

Coxeter's rabbit

On p.13 of his "Introduction to Geometry", H.S.M. Coxeter invites the reader to see (and to use spontaneously) that with  $s = (a+b+c)/2$ ,  $abc$  equals

$$(0) \quad s(s-b)(s-c) + s(s-c)(s-a) + s(s-a)(s-b) - (s-a)(s-b)(s-c)$$

Proof  $s(s-b)(s-c) + s(s-c)(s-a)$

$$= \{ \text{algebra} \}$$

$$s(s-c)(2s-a-b)$$

$$= \{ \text{definition of } s \}$$

$$(1) \quad s(s-c)c$$

$$s(s-a)(s-b) - (s-a)(s-b)(s-c)$$

$$= \{ \text{algebra} \}$$

$$(2) \quad (s-a)(s-b)c$$

Because both expressions (1) and (2) contain a factor  $c$ , so does (0); for reasons of symmetry, (0) also contains factors  $a$  and  $b$ , i.e. is a multiple of  $abc$ . The coefficient equals 1 - as is trivially established with, say,  $a, b, c := 2, 2, 2$  - and thus  $abc = (0)$  has been proved. (End of Proof)

Nuenen, 14 April 2002

prof. dr. Edsger W. Dijkstra  
 Plataanstraat 5  
 5671 AL Nuenen  
 The Netherlands