#|

Copyright (C) 1994 by J Strother Moore. All Rights Reserved.

This script is hereby placed in the public domain, and therefore unlimited editing and redistribution is permitted.

NO WARRANTY

J Strother Moore PROVIDES ABSOLUTELY NO WARRANTY. THE EVENT SCRIPT IS PROVIDED "AS IS" WITHOUT WARRANTY OF ANY KIND, EITHER EXPRESS OR IMPLIED, INCLUDING, BUT NOT LIMITED TO, ANY IMPLIED WARRANTIES OF MERCHANTABILITY AND FITNESS FOR A PARTICULAR PURPOSE. THE ENTIRE RISK AS TO THE QUALITY AND PERFORMANCE OF THE SCRIPT IS WITH YOU. SHOULD THE SCRIPT PROVE DEFECTIVE, YOU ASSUME THE COST OF ALL NECESSARY SERVICING, REPAIR OR CORRECTION.

IN NO EVENT WILL J Strother Moore BE LIABLE TO YOU FOR ANY DAMAGES, ANY LOST PROFITS, LOST MONIES, OR OTHER SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES ARISING OUT OF THE USE OR INABILITY TO USE THIS SCRIPT (INCLUDING BUT NOT LIMITED TO LOSS OF DATA OR DATA BEING RENDERED INACCURATE OR LOSSES SUSTAINED BY THIRD PARTIES), EVEN IF YOU HAVE ADVISED US OF THE POSSIBILITY OF SUCH DAMAGES, OR FOR ANY CLAIM BY ANY OTHER PARTY.''

|#

EVENT: Start with the initial **nqthm** theory.

EVENT: For efficiency, compile those definitions not yet compiled.

EVENT: Add the shell *zn*, with recognizer function symbol *znp* and 2 accessors: *pos*, with type restriction (one-of numberp) and default value zero; *neg*, with type restriction (one-of numberp) and default value zero.

DEFINITION:
$\text{zlessp}(x, y) = ((\text{pos}(x) + \text{neg}(y)) < (\text{neg}(x) + \text{pos}(y)))$

DEFINITION: $\text{zlesseqp}(x, y) = (\neg\ \text{zlessp}(y, x))$

DEFINITION:
$\text{zmax}(x, y)$
$=$ **if** $\text{zlessp}(x, y)$ **then** y
else x **endif**

DEFINITION:
zmin $(x,\ y)$
$=$ **if** zlessp $(x,\ y)$ **then** x
else y **endif**

DEFINITION: zsub1 (x) = zn (pos (x), 1 + neg (x))

DEFINITION:
pzdifference $(x,\ y)$ = ((pos (x) + neg (y)) − (neg (x) + pos (y)))

DEFINITION:
m1 $(x,\ y,\ z)$
$=$ **if** zlesseqp $(x,\ y)$ **then** 0
else 1 **endif**

DEFINITION:
m2 $(x,\ y,\ z)$ = pzdifference (zmax $(x,$ zmax $(y,\ z))$, zmin $(x,$ zmin $(y,\ z)))$

DEFINITION: m3 $(x,\ y,\ z)$ = pzdifference $(x,$ zmin $(x,$ zmin $(y,\ z)))$

DEFINITION:
tak0 $(x,\ y,\ z)$
$=$ **if** zlesseqp $(x,\ y)$ **then** y
elseif zlesseqp $(y,\ z)$ **then** z
else x **endif**

DEFINITION:
m $(x,\ y,\ z)$ = cons (m1 $(x,\ y,\ z)$, cons (m2 $(x,\ y,\ z)$, cons (m3 $(x,\ y,\ z)$, **nil**)))

THEOREM: tak0-satisfies-tak-equation
tak0 $(x,\ y,\ z)$
$=$ **if** zlesseqp $(x,\ y)$ **then** y
else tak0 (tak0 (zsub1 (x), y, z),
tak0 (zsub1 (y), z, x),
tak0 (zsub1 (z), x, y)) **endif**

THEOREM: m1-lesseqp-0
($\neg$ zlesseqp $(x,\ y)$)
$\rightarrow$ (m1 $(x,\ y,\ z)$
$\not<$ m1 (tak0 (zsub1 (x), y, z), tak0 (zsub1 (y), z, x), tak0 (zsub1 (z), x, y)))

THEOREM: m1-lesseqp-1
($\neg$ zlesseqp $(x,\ y)$) $\rightarrow$ (m1 $(x,\ y,\ z)$ $\not<$ m1 (zsub1 (x), y, z))

THEOREM: m1-lesseqp-2
($\neg$ zlesseqp $(x,\ y)$) $\rightarrow$ (m1 $(x,\ y,\ z)$ $\not<$ m1 (zsub1 (y), z, x))

THEOREM: m1-lesseqp-3
$(\neg\ \text{zlesseqp}\,(x,\ y)) \rightarrow (\text{m1}\,(x,\ y,\ z) \not< \text{m1}\,(\text{zsub1}\,(z),\ x,\ y))$

THEOREM: m2-lesseqp-0
$(\neg\ \text{zlesseqp}\,(x,\ y))$
$\rightarrow\quad (\text{m2}\,(x,\ y,\ z)$
$\quad\quad \not< \quad \text{m2}\,(\text{tak0}\,(\text{zsub1}\,(x),\ y,\ z),\ \text{tak0}\,(\text{zsub1}\,(y),\ z,\ x),\ \text{tak0}\,(\text{zsub1}\,(z),\ x,\ y)))$

THEOREM: m2-lesseqp-1
$(\neg\ \text{zlesseqp}\,(x,\ y)) \rightarrow (\text{m2}\,(x,\ y,\ z) \not< \text{m2}\,(\text{zsub1}\,(x),\ y,\ z))$

THEOREM: m2-lesseqp-2
$((\neg\ \text{zlesseqp}\,(x,\ y)) \wedge (\text{m1}\,(\text{zsub1}\,(y),\ z,\ x) = \text{m1}\,(x,\ y,\ z)))$
$\rightarrow\quad (\text{m2}\,(x,\ y,\ z) \not< \text{m2}\,(\text{zsub1}\,(y),\ z,\ x))$

THEOREM: m2-lesseqp-3
$((\neg\ \text{zlesseqp}\,(x,\ y)) \wedge (\text{m1}\,(\text{zsub1}\,(z),\ x,\ y) = \text{m1}\,(x,\ y,\ z)))$
$\rightarrow\quad (\text{m2}\,(x,\ y,\ z) \not< \text{m2}\,(\text{zsub1}\,(z),\ x,\ y))$

THEOREM: m3-lessp-0
$((\neg\ \text{zlesseqp}\,(x,\ y))$
$\wedge\quad (\text{m1}\,(\text{tak0}\,(\text{zsub1}\,(x),\ y,\ z),\ \text{tak0}\,(\text{zsub1}\,(y),\ z,\ x),\ \text{tak0}\,(\text{zsub1}\,(z),\ x,\ y))$
$\quad\quad = \quad \text{m1}\,(x,\ y,\ z)))$
$\rightarrow\quad (\text{m3}\,(\text{tak0}\,(\text{zsub1}\,(x),\ y,\ z),\ \text{tak0}\,(\text{zsub1}\,(y),\ z,\ x),\ \text{tak0}\,(\text{zsub1}\,(z),\ x,\ y))$
$\quad\quad < \quad \text{m3}\,(x,\ y,\ z))$

THEOREM: m3-lessp-1
$((\neg\ \text{zlesseqp}\,(x,\ y)) \wedge (\text{m1}\,(\text{zsub1}\,(x),\ y,\ z) = \text{m1}\,(x,\ y,\ z)))$
$\rightarrow\quad (\text{m3}\,(\text{zsub1}\,(x),\ y,\ z) < \text{m3}\,(x,\ y,\ z))$

THEOREM: m3-lessp-2
$((\neg\ \text{zlesseqp}\,(x,\ y)) \wedge (\text{m1}\,(\text{zsub1}\,(y),\ z,\ x) = \text{m1}\,(x,\ y,\ z)))$
$\rightarrow\quad (\text{m3}\,(\text{zsub1}\,(y),\ z,\ x) < \text{m3}\,(x,\ y,\ z))$

THEOREM: m3-lessp-3
$((\neg\ \text{zlesseqp}\,(x,\ y)) \wedge (\text{m1}\,(\text{zsub1}\,(z),\ x,\ y) = \text{m1}\,(x,\ y,\ z)))$
$\rightarrow\quad (\text{m2}\,(\text{zsub1}\,(z),\ x,\ y) < \text{m2}\,(x,\ y,\ z))$

EVENT: Disable zlessp.

EVENT: Disable m1.

EVENT: Disable m2.

EVENT: Disable m3.

EVENT: Disable tak0.

EVENT: Disable zsub1.

DEFINITION:
make-ordinal3 (x)
$=$ cons (cons $(1 +$ car (x), 0), cons $(1 +$ cadr (x), fix (caddr (x)))))

THEOREM: ordinalp-make-ordinal3
ordinalp (make-ordinal3 (x))

DEFINITION:
lex3 (x, y) = ord-lessp (make-ordinal3 (x), make-ordinal3 (y))

THEOREM: m-goes-down-1
$(\neg$ zlesseqp $(x, y)) \rightarrow$ lex3 (m (zsub1 (x), y, z), m (x, y, z))

THEOREM: m-goes-down-2
$(\neg$ zlesseqp $(x, y)) \rightarrow$ lex3 (m (zsub1 (y), z, x), m (x, y, z))

THEOREM: m-goes-down-3
$(\neg$ zlesseqp $(x, y)) \rightarrow$ lex3 (m (zsub1 (z), x, y), m (x, y, z))

THEOREM: m-goes-down-0
$(\neg$ zlesseqp $(x, y))$
$\rightarrow$ lex3 (m (tak0 (zsub1 (x), y, z), tak0 (zsub1 (y), z, x), tak0 (zsub1 (z), x, y)),
m (x, y, z))

Index

lex3, 4

m, 2, 4
m-goes-down-0, 4
m-goes-down-1, 4
m-goes-down-2, 4
m-goes-down-3, 4
m1, 2, 3
m1-lesseqp-0, 2
m1-lesseqp-1, 2
m1-lesseqp-2, 2
m1-lesseqp-3, 3
m2, 2, 3
m2-lesseqp-0, 3
m2-lesseqp-1, 3
m2-lesseqp-2, 3
m2-lesseqp-3, 3
m3, 2, 3
m3-lessp-0, 3
m3-lessp-1, 3
m3-lessp-2, 3
m3-lessp-3, 3
make-ordinal3, 4

neg, 1, 2

ordinalp-make-ordinal3, 4

pos, 1, 2
pzdifference, 2

tak0, 2–4
tak0-satisfies-tak-equation, 2

zlesseqp, 1–4
zlessp, 1, 2
zmax, 1, 2
zmin, 2
zn, 1, 2
zsub1, 2–4