

CS311H: Discrete Mathematics

Recurrence Relations

Instructor: Işıl Dillig

Recall: Recursively Defined Sequences

- ▶ In previous lectures, we looked at **recursively-defined sequences**
- ▶ **Example:** What sequence is this?

$$\begin{aligned}a_0 &= 1 \\a_1 &= 1 \\a_n &= a_{n-1} + a_{n-2}\end{aligned}$$

Recurrence Relations

- ▶ Recursively defined sequences are often referred to as **recurrence relations**
- ▶ The base cases in the recursive definition are called **initial values** of the recurrence relation
- ▶ **Example:** Write recurrence relation representing number of bacteria in n 'th hour if colony starts with 5 bacteria and doubles every hour?

Closed Form Solutions

- ▶ Often, we need to find a **closed form solution** for a given recurrence
- ▶ **Recall:** Closed form solution defines n 'th number in the sequence as a function of n
- ▶ What is closed form solution to the following recurrence?

$$\begin{aligned}a_0 &= 0 \\a_n &= a_{n-1} + n\end{aligned}$$

Closed Form Solutions of Recurrence Relations

- ▶ Given an arbitrary recurrence relation, is there a mechanical way to obtain the **closed form solution**?
- ▶ Not for arbitrary, but for a subclass of recurrence relations
- ▶ A **linear homogeneous** recurrence relation with constant coefficients is a recurrence relation of the form:

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \dots + c_k a_{n-k}$$

where each c_i is a constant and c_k is non-zero

- ▶ The value of k is called the **degree** of the recurrence relation

Examples and Non-Examples

- ▶ Which of these are linear homogeneous recurrence relations with constant coefficients?
 - ▶ $a_n = a_{n-1} + 2a_{n-5}$
 - ▶ $a_n = 2a_{n-2} + 5$
 - ▶ $a_n = a_{n-1} + n$
 - ▶ $a_n = a_{n-1} \cdot a_{n-2}$
 - ▶ $a_n = n \cdot a_{n-1}$

Characteristic Polynomial

- ▶ Cook-book recipe for solving linear homogenous recurrence relations with constant coefficients
- ▶ **Definition:** The **characteristic equation** of a recurrence relation of the form $a_n = c_1 a_{n-1} + c_2 a_{n-2} + \dots + c_k a_{n-k}$ is

$$r^k = c_1 r^{k-1} + c_2 r^{k-2} + \dots + c_k$$

- ▶ i.e., replace a_i with $r^{i-(n-k)}$

Characteristic Equation Examples

- ▶ What are the characteristic equations for the following recurrence relations?

- ▶ $f_n = f_{n-1} + f_{n-2}$

- ▶ $a_n = 2a_{n-1}$

- ▶ $a_n = 2a_{n-1} + 5a_{n-3}$

Characteristic Roots

- ▶ The **characteristic roots** of a linear homogeneous recurrence relation are the roots of its characteristic equation.
- ▶ What are the characteristic roots of the following recurrence relations?

- ▶ $a_n = 2a_{n-1} + 3a_{n-2}$

- ▶ $f_n = f_{n-1} + f_{n-2}$

Theorem I for Solving Linear Homogenous Recurrence Relations

Let $a_n = c_1 a_{n-1} + c_2 a_{n-2} + \dots + c_k a_{n-k}$ be a recurrence relation with k **distinct** characteristic roots r_1, \dots, r_k .

- ▶ Then the closed form solution for a_n is of the form:

$$\alpha_1 r_1^n + \alpha_2 r_2^n + \dots + \alpha_k r_k^n$$

- ▶ Furthermore, given k initial conditions, the constants $\alpha_1, \dots, \alpha_k$ are **uniquely determined**
- ▶ **Note:** Won't do the proof because requires a good amount of linear algebra

Example

Find a closed form solution for the recurrence $a_n = a_{n-1} + 2a_{n-2}$ with initial conditions $a_0 = 2$ and $a_1 = 7$

- ▶ **Characteristic equation:**
- ▶ **Characteristic roots:**
- ▶ **Coefficients:**
- ▶ **Closed-form solution:**

Generalized Theorem

- ▶ So far, we assume all characteristic roots are **distinct** – what happens if this is not the case?
- ▶ **Theorem:** Let $a_n = c_1 a_{n-1} + c_2 a_{n-2} + \dots + c_k a_{n-k}$ be a recurrence relation with t distinct characteristic roots r_1, \dots, r_k with **multiplicities** m_1, \dots, m_k . Then solutions are of the form:

$$a_n = \sum_{i=0}^t (\alpha_{i,0} + \alpha_{i,1} \cdot n + \dots + \alpha_{i,m_i-1} \cdot n^{m_i-1}) r_i^n$$

An Example

- ▶ Find closed form of recurrence $a_n = 3a_{n-1} - 3a_{n-2} + a_{n-3}$ with initial conditions $a_0 = 1, a_1 = 3, a_2 = 7$
- ▶ Characteristic equation:
- ▶ Characteristic roots:
- ▶ Solution form:
- ▶ Coefficients:

Solving Linear Non-Homogeneous Recurrence Relations

- ▶ How do we solve linear, but non-homogeneous recurrence relations, such as $a_n = 2a_{n-1} + 1$?
- ▶ A **linear non-homogeneous** recurrence relation with constant coefficients is of the form:

$$a_n = c_1 a_{n-1} + a_2 a_{n-2} + \dots + c_k a_{n-k} + F(n)$$

- ▶ The recurrence obtained by dropping $F(n)$ is called the **associated homogeneous recurrence relation**
- ▶ To solve these recurrences, we will combine the solution for the homogenous recurrence with **particular solution**

Particular Solution

- ▶ A **particular solution** for a recurrence relation is one that satisfies the recurrence but not necessarily the initial conditions
- ▶ **Example:** Consider the recurrence $a_n = a_{n-1} + 1$ with initial condition $a_0 = 5$
- ▶ A particular solution for this recurrence is $a_n = n$, but it does not satisfy the initial condition

Theorem about Linear Non-homogeneous Recurrences

Suppose $a_n = c_1 a_{n-1} + \dots + c_k a_{n-k} + F(n)$ has **particular solution** a_n^p , and a_n^h is solution for associated homogeneous recurrence. Then every solution is of the form $a_n^p + a_n^h$.

Why is this theorem useful?

- ▶ If we can find a particular solution, then we can also mechanically find a solution that satisfies initial conditions.
- ▶ **Example:** Solve the recurrence relation $a_n = 3a_{n-1} + 2n$ with initial condition $a_1 = 3$
- ▶ A **particular solution:** $-n - \frac{3}{2}$ (Why?)
- ▶ Solutions for homogeneous recurrence:
- ▶ Solutions for recurrence:
- ▶ Solve for α :

How do we find a particular solution?

Theorem: Consider $a_n = c_1 a_{n-1} + \dots + c_k a_{n-k} + F(n)$ where:

$$F(n) = (b_t n^t + b_{t-1} n^{t-1} + \dots + b_1 n + b_0) s^n$$

- ▶ **Case 1:** If s is not a root of the associated characteristic equation, then there exists a **particular solution** of the form:

$$(p_t n^t + p_{t-1} n^{t-1} + \dots + p_1 n + p_0) s^n$$

- ▶ **Case 2:** If s is a root with multiplicity m of the characteristic equation, then there exists a solution of the form:

$$n^m (p_t n^t + p_{t-1} n^{t-1} + \dots + p_1 n + p_0) s^n$$

Example I

- ▶ Consider again the recurrence $a_n = 3a_{n-1} + 2n$
- ▶ Here, $s = 1$ and characteristic root is 3
- ▶ Hence, there exists a particular solution of the form $p_1 n + p_0$
- ▶ Now, solve for p_0, p_1 :


$$p_1 n + p_0 = 3(p_1(n-1) + p_0) + 2n$$

- ▶ Rearrange: $2n(p_1 + 1) + (2p_0 - 3p_1) = 0$
- ▶ A solution $p_1 = -1, p_0 = -\frac{3}{2}$
- ▶ A particular solution: $-n - \frac{3}{2}$

Example II

- ▶ Find a particular solution for $a_n = 6a_{n-1} - 9a_{n-2} + 2^n$
- ▶ **Characteristic root:**
- ▶ Particular solution of the form:
- ▶ Find p_0 such that $p_0 \cdot 2^n = 6(p_0 \cdot 2^{n-1}) - 9(p_0 \cdot 2^{n-2}) + 2^n$
- ▶ Solve for p_0 :
- ▶ Particular solution:

Towers of Hanoi



- ▶ Given 3 pegs where first peg contains n disks
- ▶ **Goal:** Move all the disks to a different peg (e.g., second one)
- ▶ **Rule 1:** Larger disks cannot rest on top of smaller disks
- ▶ **Rule 2:** Can only move the top-most disk at a time
- ▶ **Question:** How many steps does it take to move all n disks?

A Recursive Solution

- ▶ Solve recursively – T_n is number of steps to move n disks
- ▶ **Base case:** $n = 1$, move disk from first peg to second: $T_1 = 1$
- ▶ **Induction:** Suppose we can move $n - 1$ disks in T_{n-1} steps; how many steps does it take to move T_n disks?
- ▶ **Idea:** First move the topmost $n - 1$ disks to peg 3; can be done in T_{n-1} steps
- ▶ Now, move bottom-most disk to peg 2 – takes just 1 step
- ▶ Finally, recursively move $n - 1$ disks in peg 3 to peg 2 – can be done in T_{n-1} steps

Towers of Hanoi, cont.

- ▶ **Recurrence relation:**
- ▶ **Initial condition:**
- ▶ Now find closed form for T_n
- ▶ What is a particular solution?
- ▶ Solution for homogeneous recurrence:
- ▶ Solve for α :
- ▶ Solution for recurrence: