

CS311H: Discrete Mathematics

Combinatorics

Instructor: Işıl Dillig

Combinatorics

- ▶ **Combinatorics** (counting) deals with the question: "How many elements in a given set have desired property?"
- ▶ Counting problems can be hard \Rightarrow useful to decompose
- ▶ Two basic very useful decomposition rules:
 1. **Product rule**: useful when task decomposes into a sequence of independent tasks
 2. **Sum rule**: decomposes task into a set of alternatives

Product Rule

- ▶ Suppose a task A can be decomposed into a sequence of two independent tasks B and C
- ▶ n_1 ways of doing B
- ▶ n_2 ways of doing C
- ▶ **Product rule**: Then, there are $n_1 n_2$ ways of doing A

Example 1

- ▶ New company with 12 offices and 2 employees Kate and Jack
- ▶ How many ways to assign different offices to Kate and Jack?
- ▶ **Decomposition**: First assign office to Kate, then to Jack
- ▶
- ▶
- ▶

Example 2

- ▶ Chairs in auditorium labeled with a letter (A-Z) and an integer $\in [1, 100]$.
- ▶ What is the max number of chairs that can be labeled?
- ▶ **Observe**: Max # of labeled chairs = # of different labelings
- ▶ **Decomposition**: First assign letter, then integer to chair
- ▶
- ▶

Example 3: Extended Product Rule

- ▶ Product rule generalizes to any k tasks
- ▶ If there are n_1 ways of doing A_1, \dots, n_k ways of doing A_k , then there are $n_1 \cdot n_2 \cdot \dots \cdot n_k$ ways of doing A
- ▶ A **bitstring** is a string where each character is either 0 or 1
- ▶ How many different bit strings of length 7 are there?
- ▶

Counting One-to-One Functions

- ▶ How many **one-to-one** functions are there from a set with 3 elements to a set with 5 elements?
- ▶
- ▶
- ▶
- ▶
- ▶

Sum Rule

- ▶ Counting problems can be hard \Rightarrow useful to decompose
- ▶ Two basic very useful decomposition rules:
 1. **Product rule** ✓
 2. **Sum rule**
- ▶ Suppose a task A can be done **either** in way B or in way C
- ▶ Suppose there are n_1 ways to do B , and n_2 ways to do C
- ▶ **Sum rule:** There are $n_1 + n_2$ ways to do A .

Example 1

- ▶ Suppose either a CS faculty or CS student must be chosen as representative for a committee
- ▶ There are 14 faculty, and 50 majors
- ▶ How many ways are there to choose the representative?
- ▶ By the sum rule, $50 + 14 = 64$ ways
- ▶ **Note:** Just like the product rule, the sum rule can be extended to more than two tasks

Example 2

- ▶ A student can choose a senior project from one of three lists
- ▶ First list contains 23 projects; second list has 15 projects, and third has 19 projects
- ▶ Also, no project appears on more than one list
- ▶ How many different projects can student choose?
- ▶
- ▶ What if some of the projects appeared on both lists?
- ▶ **Caveat:** For sum rule to apply, the possibilities must be **mutually exclusive**

More Complex Counting Problems

- ▶ Problems so far required either only product or only sum rule
- ▶ But more complex problems require a combination of both!
- ▶ **Example:** In a programming language, a variable name is a string of one or two characters.
- ▶ A character is either a letter $[a-z]$ or a digit $[0,9]$, and first character must be a letter.
- ▶ How many possible variable names are there?

Example, cont.

Another Example

- ▶ A password must be **six to seven** characters long
- ▶ A character is upper case letter or digit
- ▶ Each password must contain **at least one digit**
- ▶ How many possible passwords?
- ▶
- ▶

Example, cont.

Example 3

- ▶ How many bitstrings are there of length 6 that do not have **two consecutive 1's**?
- ▶ Let $F(n)$ denote the number of bitstrings of length n that do not have two consecutive 1's
- ▶ We'll first derive a recursive equation to characterize $F(n)$
- ▶ By the **sum rule**, $F(n)$ is the sum of:
 1. # of n -bit strings starting with 1 not containing 11
 2. # of n -bit strings starting with 0 not containing 11

Example 3, cont.


Recall: Sum Rule

- ▶ **Recall:** Sum rule only applies if a task is as disjunction of two mutually exclusive tasks
- ▶ What do we do if the tasks aren't mutually exclusive?
- ▶ **Example:** You can choose from set A or set B , but they have some elements in common
- ▶ Generalization of the sum rule: **inclusion-exclusion principle**

The Inclusion-Exclusion Principle

- ▶ Suppose a set A can be written as union of sets B and C
- ▶ **Inclusion-Exclusion Principle:**

$$|A| = |B| + |C| - |B \cap C|$$



Inclusion-Exclusion Principle Example

- ▶ How many bit strings of length 8 either start with 1 or end with two bits 00?
- ▶ Let B be the set of bitstrings that start with 1
- ▶ Let C be the set of bitstrings that end with 00
- ▶ We want $|B \cup C|$
- ▶ By the inclusion-exclusion principle,
 $|B \cup C| = |B| + |C| - |B \cap C|$
- ▶ Thus, compute $|B|$, $|C|$ and $|B \cap C|$

Example, cont.

Another Example

- ▶ A company receives 350 applications for job positions
- ▶ 220 of applicants are CS majors
- ▶ 147 of applicants are business majors
- ▶ 51 are double CS and business majors
- ▶ How many are neither CS nor business majors?
- ▶
- ▶
- ▶

The Pigeonhole Principle



- ▶ Suppose there is a flock of 36 pigeons and a set of 35 pigeonholes
- ▶ Each pigeon want to sit in one hole
- ▶ But since there are less holes than there are pigeons, one pigeon is left without a hole.

- ▶ **The Pigeonhole Principle:** If $n + 1$ or more objects are placed into n boxes, then at least one box contains 2 or more objects

Examples

- ▶ Consider an event with 367 people. Is it possible no pair of people have the same birthday?
- ▶ Consider function f from a set with $k + 1$ or more elements to a set with k elements. Is it possible f is one-to-one?
- ▶ Consider n married couples. How many of the $2n$ people must be selected to guarantee there is at least one married couple?

Generalized Pigeonhole Principle

- ▶ If n objects are placed into k boxes, then there is at least one box containing at least $\lceil n/k \rceil$ objects
- ▶ **Proof:** (by contradiction) Suppose every box contains less than $\lceil n/k \rceil$ objects
- ▶
- ▶
- ▶
- ▶

Examples

- ▶ If there are 30 students in a class, at least how many must be born in the same month? $\lceil \frac{30}{12} \rceil = 3$
- ▶ What is the minimum # of students required to ensure at least 6 students receive the same grade (A, B, C, D, F)?
- ▶
- ▶ What is the min # of cards that must be chosen to guarantee three have same suit?
- ▶