

CS311H: Discrete Mathematics

Introduction to First-Order Logic

Instructor: Işıl Dillig

Why First-Order Logic?

- ▶ So far, we studied the simplest logic: **propositional logic**
- ▶ But for some applications, propositional logic is not expressive enough
- ▶ First-order logic is more expressive: allows representing more complex facts and making more sophisticated inferences

A Motivating Example

- ▶ For instance, consider the statement “Anyone who drives fast gets a speeding ticket”
- ▶ From this, we should be able to conclude “If Joe drives fast, he will get a speeding ticket”
- ▶ Similarly, we should be able to conclude “If Rachel drives fast, she will get a speeding ticket” and so on.
- ▶ But PL does not allow inferences like that because we cannot talk about concepts like “everyone”, “someone” etc.
- ▶ **First-order logic** (predicate logic) allows making such kinds of inferences

Building Blocks of First-Order Logic

- ▶ The building blocks of propositional logic were **propositions**
- ▶ In first-order logic, there are three kinds of basic building blocks: constants, variables, predicates
- ▶ **Constants**: refer to specific objects (in a universe of discourse)
- ▶ **Examples**: George, 6, Austin, CS311, ...
- ▶ **Variables**: range over objects (in a universe of discourse)
- ▶ **Examples**: x, y, z , ...
- ▶ If universe of discourse is cities in Texas, x can represent Houston, Austin, Dallas, San Antonio, ...

Building Blocks of First-Order Logic, cont.

- ▶ **Predicates** describe properties of objects or relationships between objects
- ▶ **Examples:** ishappy, betterthan, loves, $>$. . .
- ▶ Predicates can be applied to both constants and variables
- ▶ **Examples:** ishappy(George), betterthan(x,y), loves(George, Rachel), $x > 3$, . . .
- ▶ A predicate $P(c)$ is true or false depending on whether property P holds for c
- ▶ **Example:** ishappy(George) is true if George is happy, but false otherwise

Predicate Examples

- ▶ Consider predicate **even** which represents if a number is even
- ▶ What is truth value of **even(2)**?
- ▶ What is truth value of **even(5)**?
- ▶ What is truth value of **even(x)**?
- ▶ Another example: Suppose $Q(x, y)$ denotes $x = y + 3$
- ▶ What is the truth value of $Q(3, 0)$?
- ▶ What is the truth value of $Q(1, 2)$?

Formulas in First Order Logic

- ▶ Formulas in first-order logic are formed using predicates and logical connectives.
- ▶ Example: $\text{even}(2)$ is a formula
- ▶ Example: $\text{even}(x)$ is also a formula
- ▶ Example: $\text{even}(x) \vee \text{odd}(x)$ is also a formula
- ▶ Example: $(\text{odd}(x) \rightarrow \neg \text{even}(x)) \wedge \text{even}(x)$

Semantics of First-Order Logic

- ▶ In propositional logic, the truth value of formula depends on a truth assignment to variables.
- ▶ In FOL, truth value of a formula depends **interpretation** of predicate symbols and variables over some domain D
- ▶ Consider a FOL formula $\neg P(x)$
- ▶ A possible interpretation:

$$D = \{\star, \circ\}, P(\star) = \text{true}, P(\circ) = \text{false}, x = \star$$

- ▶ Under this interpretation, what's truth value of $\neg P(x)$?
- ▶ What about if $x = \circ$?

More Examples

- ▶ Consider interpretation I over domain $D = \{1, 2\}$
 - ▶ $P(1, 1) = P(1, 2) = \text{true}$, $P(2, 1) = P(2, 2) = \text{false}$
 - ▶ $Q(1) = \text{false}$, $Q(2) = \text{true}$
 - ▶ $x = 1$, $y = 2$
- ▶ What is truth value of $P(x, y) \wedge Q(y)$ under I ?
- ▶ What is truth value of $P(y, x) \rightarrow Q(y)$ under I ?
- ▶ What is truth value of $P(x, y) \rightarrow Q(x)$ under I ?

Quantifiers

- ▶ Real power of first-order logic over propositional logic:
quantifiers
- ▶ Quantifiers allow us to talk about **all** objects or the existence of **some** object
- ▶ There are two quantifiers in first-order logic:
 1. Universal quantifier ($\forall$): refers to **all** objects
 2. Existential quantifier ($\exists$): refers to **some** object

Universal Quantifiers

- ▶ **Universal quantification** of $P(x)$, $\forall x.P(x)$, is the statement "P(x) holds for all objects x in the universe of discourse."
- ▶ $\forall x.P(x)$ is true if predicate P is true for **every** object in the universe of discourse, and false otherwise
- ▶ Consider domain $D = \{\circ, \star\}$, $P(\circ) = \text{true}$, $P(\star) = \text{false}$
- ▶ What is truth value of $\forall x.P(x)$?
- ▶ Object $\circ$ for which $P(\circ)$ is false is **counterexample** of $\forall x.P(x)$
- ▶ What is a counterexample for $\forall x.P(x)$ in previous example?

More Universal Quantifier Examples

- ▶ Consider the domain D of real numbers and predicate $P(x)$ with interpretation $x^2 \geq x$
- ▶ What is the truth value of $\forall x.P(x)$?
- ▶ What is a counterexample?
- ▶ What if the domain is integers?
- ▶ **Observe:** Truth value of a formula depends on a universe of discourse!

Existential Quantifiers

- ▶ **Existential quantification** of $P(x)$, written $\exists x.P(x)$, is "There exists an element x in the domain such that $P(x)$ ".
- ▶ $\exists x.P(x)$ is true if there is **at least one** element in the domain such that $P(x)$ is true
- ▶ In first-order logic, domain is required to be **non-empty**.
- ▶ Consider domain $D = \{\circ, \star\}$, $P(\circ) = \text{true}$, $P(\star) = \text{false}$
- ▶ What is truth value of $\exists x.P(x)$?

Existential Quantifier Examples

- ▶ Consider the domain of reals and predicate $P(x)$ with interpretation $x < 0$.
- ▶ What is the truth value of $\exists x.P(x)$?
- ▶ What if domain is positive integers?
- ▶ Let $Q(y)$ be the statement $y > y^2$
- ▶ What's truth value of $\exists y.Q(y)$ if domain is reals?
- ▶ What about if domain is integers?

Quantifiers Summary

Statement	When True?	When False?
$\forall x.P(x)$	$P(x)$ is true for every x	$P(x)$ is false for some x
$\exists x.P(x)$	$P(x)$ is true for some x	$P(x)$ is false for every x

- ▶ Consider finite universe of discourse with objects $o_1, \dots, o_n$
- ▶ $\forall x.P(x)$ is true iff $P(o_1) \wedge P(o_2) \dots \wedge P(o_n)$ is true
- ▶ $\exists x.P(x)$ is true iff $P(o_1) \vee P(o_2) \dots \vee P(o_n)$ is true

Quantified Formulas

- ▶ So far, only discussed how to quantify individual predicates.
- ▶ But we can also quantify entire formulas containing multiple predicates and logical connectives.
- ▶ $\exists x.(\text{even}(x) \wedge \text{gt}(x, 100))$ is a valid formula in FOL
- ▶ What's truth value of this formula if domain is all integers?
 - ▶ assuming $\text{even}(x)$ means " x is even" and $\text{gt}(x, y)$ means $x > y$
- ▶ What about $\forall x.(\text{even}(x) \wedge \text{gt}(x, 100))$?

More Examples of Quantified Formulas

- ▶ Consider the domain of integers and the predicates $\text{even}(x)$ and $\text{div4}(x)$ which represents if x is divisible by 4
- ▶ What is the truth value of the following quantified formulas?
 - ▶ $\forall x. (\text{div4}(x) \rightarrow \text{even}(x))$
 - ▶ $\forall x. (\text{even}(x) \rightarrow \text{div4}(x))$
 - ▶ $\exists x. (\neg \text{div4}(x) \wedge \text{even}(x))$
 - ▶ $\exists x. (\neg \text{div4}(x) \rightarrow \text{even}(x))$
 - ▶ $\forall x. (\neg \text{div4}(x) \rightarrow \text{even}(x))$

Translating English Into Quantified Formulas

Assuming $\text{freshman}(x)$ means “ x is a freshman” and $\text{inCS311}(x)$ “ x is taking CS311”, express the following in FOL

- ▶ Someone in CS311 is a freshman
- ▶ No one in CS311 is a freshman
- ▶ Everyone taking CS311 are freshmen
- ▶ Every freshman is taking CS311

DeMorgan's Laws for Quantifiers

- ▶ Learned about DeMorgan's laws for propositional logic:

$$\begin{aligned}\neg(p \wedge q) &\equiv \neg p \vee \neg q \\ \neg(p \vee q) &\equiv \neg p \wedge \neg q\end{aligned}$$

- ▶ DeMorgan's laws extend to first-order logic, e.g.,
 $\neg(\text{even}(x) \vee \text{div4}(x)) \equiv (\neg\text{even}(x) \wedge \neg\text{div4}(x))$

- ▶ Two new DeMorgan's laws for quantifiers:

$$\begin{aligned}\neg\forall x.P(x) &\equiv \exists x.\neg P(x) \\ \neg\exists x.P(x) &\equiv \forall x.\neg P(x)\end{aligned}$$

- ▶ When you push negation in, $\forall$ flips to $\exists$ and vice versa

Using DeMorgan's Laws

- ▶ Expressed "Noone in CS311 is a freshman" as
 $\neg \exists x. (\text{inCS311}(x) \wedge \text{freshman}(x))$
- ▶ Let's apply DeMorgan's law to this formula:
- ▶ Using the fact that $p \rightarrow q$ is equivalent to $\neg p \vee q$, we can write this formula as:
- ▶ Therefore, these two formulas are equivalent!

Nested Quantifiers

- ▶ Sometimes may be necessary to use multiple quantifiers
- ▶ For example, can't express "Everybody loves someone" using a single quantifier
- ▶ Suppose predicate $\text{loves}(x, y)$ means "Person x loves person y "
- ▶ What does $\forall x. \exists y. \text{loves}(x, y)$ mean?
- ▶ What does $\exists y. \forall x. \text{loves}(x, y)$ mean?
- ▶ **Observe:** Order of quantifiers is **very** important!

More Nested Quantifier Examples

Using the $\text{loves}(x,y)$ predicate, how can we say the following?


- ▶ "Someone loves everyone"
- ▶ "There is someone who doesn't love anyone"
- ▶ "There is someone who is not loved by anyone"
- ▶ "Everyone loves everyone"
- ▶ "There is someone who doesn't love herself/himself."

Summary of Nested Quantifiers

Statement	When True?
$\forall x.\forall y.P(x, y)$ $\forall y.\forall x.P(x, y)$	$P(x, y)$ is true for every pair x, y
$\forall x.\exists y.P(x, y)$	For every x , there is a y for which $P(x, y)$ is true
$\exists x.\forall y.P(x, y)$	There is an x for which $P(x, y)$ is true for every y
$\exists x.\exists y.P(x, y)$ $\exists y.\exists x.P(x, y)$	There is a pair x, y for which $P(x, y)$ is true

Observe: Order of quantifiers is only important if quantifiers of different kinds!


Understanding Quantifiers



Which formulas are true/false? If false, give a counterexample

- ▶ $\forall x. \exists y. (\text{sameShape}(x, y) \wedge \text{differentColor}(x, y))$
- ▶ $\forall x. \exists y. (\text{sameColor}(x, y) \wedge \text{differentShape}(x, y))$
- ▶ $\forall x. (\text{triangle}(x) \rightarrow (\exists y. (\text{circle}(y) \wedge \text{sameColor}(x, y))))$

Understanding Quantifiers, cont.



Which formulas are true/false? If false, give a counterexample

- ▶ $\forall x. \forall y. ((\text{triangle}(x) \wedge \text{square}(y)) \rightarrow \text{sameColor}(x, y))$
- ▶ $\exists x. \forall y. \neg \text{sameShape}(x, y)$
- ▶ $\forall x. (\text{circle}(x) \rightarrow (\exists y. (\neg \text{circle}(y) \wedge \text{sameColor}(x, y))))$

Translating First-Order Logic into English

Given predicates *student*(x), *atUT*(x), and *friends*(x, y), what do the following formulas say in English?

- ▶ $\forall x. ((\text{atUT}(x) \wedge \text{student}(x)) \rightarrow (\exists y. (\text{friends}(x, y) \wedge \neg \text{atUT}(y))))$
- ▶ $\forall x. ((\text{student}(x) \wedge \neg \text{atUT}(x)) \rightarrow \neg \exists y. \text{friends}(x, y))$
- ▶ $\forall x. \forall y. ((\text{student}(x) \wedge \text{student}(y) \wedge \text{friends}(x, y)) \rightarrow (\text{atUT}(x) \wedge \text{atUT}(y)))$

Translating English into First-Order Logic

Given predicates $student(x)$, $atUT(x)$, and $friends(x, y)$, how do we express the following in first-order logic?

- ▶ "Every UT student has a friend"
- ▶ "At least one UT student has no friends"
- ▶ "All UT students are friends with each other"

Satisfiability, Validity in FOL

- ▶ The concepts of satisfiability, validity also important in FOL
- ▶ An FOL formula F is satisfiable if there exists some domain and some interpretation such that F evaluates to true
- ▶ Example: Prove that $\forall x.P(x) \rightarrow Q(x)$ is satisfiable.
- ▶ An FOL formula F is valid if, for all domains and all interpretations, F evaluates to true
- ▶ Prove that $\forall x.P(x) \rightarrow Q(x)$ is not valid.
- ▶ Formulas that are satisfiable, but not valid are **contingent**, e.g., $\forall x.P(x) \rightarrow Q(x)$

Equivalence

- ▶ Two formulas F_1 and F_2 are equivalent if $F_1 \leftrightarrow F_2$ is valid
- ▶ In PL, we could prove equivalence using truth tables, but not possible in FOL
- ▶ However, we can still use known equivalences to rewrite one formula as the other
- ▶ **Example:** Prove that $\neg(\forall x. (P(x) \rightarrow Q(x)))$ and $\exists x. (P(x) \wedge \neg Q(x))$ are equivalent.
- ▶ **Example:** Prove that $\neg\exists x.\forall y.P(x, y)$ and $\forall x.\exists y.\neg P(x, y)$ are equivalent.