

Arithmetic complexity via effective names for random sequences

BJØRN KJOS-HANSEN, University of Hawai'i at Mānoa

FRANK STEPHAN, National University of Singapore

JASON R. TEUTSCH, Ruprecht-Karls-Universität Heidelberg

We investigate enumerability properties for classes of sets which permit recursive, lexicographically increasing approximations, or *left-r.e.* sets. In addition to pinpointing the complexity of left-r.e. Martin-Löf, computably, Schnorr, and Kurtz random sets, weakly 1-generics and their complementary classes, we find that there exist characterizations of the third and fourth levels of the arithmetic hierarchy purely in terms of these notions. More generally, there exists an equivalence between arithmetic complexity and existence of numberings for classes of left-r.e. sets with shift-persistent elements. While some classes (such as Martin-Löf randoms and Kurtz non-randoms) have left-r.e. numberings, there is no canonical, or *acceptable*, left-r.e. numbering for any class of left-r.e. randoms. Finally, we note some fundamental differences between left-r.e. numberings for sets and reals.

Categories and Subject Descriptors: F.4.1 [Mathematical Logic and Formal Languages]: Mathematical Logic — Computability Theory, Computational Logic

General Terms: Algorithms, Theory

Additional Key Words and Phrases: arithmetical hierarchy, Schnorr randomness, computable randomness

ACM Reference Format:

Kjos-Hanssen, B., Stephan, F., and Teutsch, J. 2012. Arithmetic complexity via effective names for random sequences. *ACM Trans. Comput. Logic* 13, 3, Article 1 (July 2012), 19 pages.

DOI = 10.1145/0000000.0000000 <http://doi.acm.org/10.1145/0000000.0000000>

1. EFFECTIVE RANDOMNESS

Think of a real number between 0 and 1. Is it random? In order to give a meaningful answer to this question, one must first obtain an expression for the real number in mind. Any reasonable language contains no more than countably many expressions, and therefore we must always settle for a language with uncountably many indescribable reals. On the other hand, there exists a natural and robust class of real numbers which admit recursive increasing approximations. We call such numbers *left-r.e.* reals. Brodhead and Kjos-Hanssen [Brodhead and Kjos-Hanssen 2009] observed that there exists an effective enumeration, or *numbering*, of the left-r.e. reals, and Chaitin [Chaitin 1987] showed that some left-r.e. reals are Martin-Löf random. Random left-r.e. reals thus serve as a friction point between definability and pure randomness.

This material is based upon work supported by the National Science Foundation under Grants No. 0652669 and 0901020 (B. Kjos-Hanssen). F. Stephan is partially supported by NUS grant R252-000-420-112. J. Teutsch is supported by the Deutsche Forschungsgemeinschaft grant ME 1806/3-1. This work is supported by the National Science Foundation, under grant CNS-0435060, grant CCR-0325197 and grant EN-CS-0329609.

Author's addresses: B. Kjos-Hanssen, Department of Mathematics, University of Hawai'i at Mānoa; F. Stephan, Departments of Mathematics and Computer Science, National University of Singapore; J. Teutsch, Department of Computer Science, Ruprecht-Karls-Universität Heidelberg.

Permission to make digital or hard copies of part or all of this work for personal or classroom use is granted without fee provided that copies are not made or distributed for profit or commercial advantage and that copies show this notice on the first page or initial screen of a display along with the full citation. Copyrights for components of this work owned by others than ACM must be honored. Abstracting with credit is permitted. To copy otherwise, to republish, to post on servers, to redistribute to lists, or to use any component of this work in other works requires prior specific permission and/or a fee. Permissions may be requested from Publications Dept., ACM, Inc., 2 Penn Plaza, Suite 701, New York, NY 10121-0701 USA, fax +1 (212) 869-0481, or permissions@acm.org.

© 2012 ACM 1529-3785/2012/07-ART1 \$10.00

DOI 10.1145/0000000.0000000 <http://doi.acm.org/10.1145/0000000.0000000>

In the following exposition we examine which classes of left-r.e. randoms and non-randoms admit numberings (and are therefore describable). A related definability question also arises, namely *how difficult is it to determine whether a real is random?* As a means of classifying complexity, we place the index sets for left-r.e. randoms inside the arithmetic hierarchy. One can view this program as a continuation of work by Hitchcock, Lutz, and Terwijn [Hitchcock et al. 2007] which places classes of randoms inside the broader Borel hierarchy. In contrast with the case of r.e. sets, we shall find a close connection between numberings and arithmetic complexity for classes of left-r.e. reals.

Notation. Some standard notation used in this article includes \forall^∞ which denotes “for all but finitely many” and \exists^∞ which means “there exist infinitely many.” $X \upharpoonright n$ is the length n prefix of X , and \cdot denotes concatenation. For finite sequences σ and τ , $\sigma \preceq \tau$ means that σ is a prefix of τ , $\sigma \prec \tau$ indicates that σ is a proper prefix of τ , and $|\sigma|$ is the length of σ . For non-negative integers x , $\lfloor x \rfloor$ is the floor of $\log(x + 1)$. $\langle \cdot, \cdot \rangle : \omega \times \omega \mapsto \omega$ is some recursive pairing function which we fix for rest of the paper. For sets A and B , $A \oplus B = \{2n : n \in A\} \cup \{2n + 1 : n \in B\}$. $'$ is the jump operator, μ is the unbounded search operator, \downarrow denotes convergence, and $A \leq_T B$ means A Turing reduces to B . As usual, \emptyset' denotes the halting set, and \bar{A} denotes the complement of the set A . For further background on recursion theory and algorithmic randomness, see [Soare 1987] and [Downey and Hirschfeldt 2010].

A *sequence* is the characteristic function of a set of natural numbers, and each sequence A corresponds to a unique real number

$$\text{real}(A) = \sum_n 2^{-n-1} \cdot A(n).$$

We denote the class of all sequences by $\{0, 1\}^\omega$, and $\{0, 1\}^*$ is the class of finite strings. A partial recursive function (synonymously, a *machine*) M is said to be *prefix-free* if for any finite strings $\sigma, \tau \in \text{dom } M$, σ is not a proper prefix of τ . The *prefix-free complexity* of a string σ with respect to a prefix-free machine M is given by $K_M(\sigma) = \min\{|p| : M(p) = \sigma\}$. Furthermore, there exists a *universal* prefix-free machine U such that for any prefix-free machine M , $K_U(\sigma) \leq K_M(\sigma) + O(1)$ for all $\sigma \in \{0, 1\}^*$ [Li and Vitányi 2008]. We fix such a U and let $K = K_U$ for the remainder of this exposition.

Definition 1.1. A sequence X is called *Martin-Löf random* [Levin 1974; Martin-Löf 1966] if

$$(\exists c) (\forall n) [K(X \upharpoonright n) \geq n - c]. \quad (1)$$

Intuitively, every prefix of the string X in (1) is incompressible and therefore admits no simple description.

A *martingale* $M : \{0, 1\}^* \rightarrow \mathbb{R} \cap [0, \infty)$ is a function satisfying the fairness condition: for all $\sigma \in 2^{<\omega}$,

$$M(\sigma) = \frac{M(\sigma 0) + M(\sigma 1)}{2}.$$

The martingale M *succeeds* on a sequence X if $\limsup M(X \upharpoonright n) = \infty$. If M succeeds on X and there exists a recursive, non-decreasing, unbounded function g satisfying $g(n) \leq M(X \upharpoonright n)$ for infinitely many n , we say that M *Schnorr-succeeds* on X . A martingale M *Kurtz-succeeds* on a set A if M succeeds on A and there exists a recursive, non-decreasing, unbounded function f such that $M(A \upharpoonright n) > f(n)$ for all n . The idea behind Definition 1.2 is that no gambling strategy can achieve arbitrary wealth by betting on a random sequence.

Definition 1.2. A sequence X is called *computably random* [Schnorr 1971a,b] if no recursive martingale succeeds on X , *Schnorr random* [Schnorr 1971b] if no recursive martingale Schnorr-succeeds on X , and *Kurtz random* [Downey et al. 2004; Kurtz 1981; Wang 1996] if no recursive martingale Kurtz-succeeds on X .

The classes of randoms mentioned above relate to each other as follows:

THEOREM 1.3 (SEE [DOWNEY AND HIRSCHFELDT 2010] OR [NIES 2009]).
Martin-Löf randomness \implies computable randomness \implies Schnorr randomness \implies Kurtz randomness.

Our discussion will also involve a related class of sequences which we introduce in Definition 1.4. A set of finite strings S is called *dense* if for every string σ there exists $\tau \in S$ extending σ .

Definition 1.4. A sequence is *weakly 1-generic* if it has a prefix in every dense r.e. set of strings. Even stronger, a sequence X is *1-generic* if for every r.e. set of strings A , either X has a prefix in A or there exists a prefix of X which has no extension in A .

While a left-r.e. real cannot be 1-generic [Odifreddi 1999, Proposition XI.2.3], weakly 1-generic sets can be left-r.e. [Jain et al. 2011; Nies 2009; Stephan and Teutsch Stephan and Teutsch]. We shall make use of the following result of Kurtz which also appears in [Downey and Hirschfeldt 2010, Theorem 8.11.7].

THEOREM 1.5 (KURTZ [1981]). *Every weakly 1-generic is Kurtz random.*

2. SETS, REALS, AND ACCEPTABLE NUMBERINGS

We turn our attention to the magical correspondence between sequences of natural numbers and reals in $[0, 1]$. In particular, the characteristic function of each subset is the binary expansion of some real number and vice-versa. We call a real *non-dyadic* if its binary expansion contains both infinitely many 1's and infinitely many 0's, and *dyadic* otherwise. This definition highlights an important distinction between sets and reals. For any string σ , the real number $.\sigma 01111\dots$ equals $.\sigma 100000\dots$. Hence there is no difference between the set of “finite” reals and the set of “co-finite” reals. For the same reason, and unlike the case for sequences, there is no difference between “infinite” and “co-infinite” reals. We shall use $\text{real}(A)$ to denote the unique real representation of a set A and $\text{set}(X)$ to denote an arbitrarily selected set representation of a real X .

In general, enumerability will depend on whether we view our objects of study as sequences or as reals, see Remark 2.4. Indeed, sequence enumerations are more restrictive than real enumerations. For random objects, however, the choice of sequences versus reals is immaterial since random reals are non-dyadic. Every random real corresponds to a unique random sequence (which in turn corresponds uniquely to the characteristic function of a set) and vice-versa. The identification of finite and co-finite sets leads to ambiguity in terminology and reference, hence we favor sets over reals throughout this exposition. Nevertheless, we keep in mind the correspondence between sets and reals and occasionally exploit their relationship. Where the discussion does not benefit from distinction between random reals, random sets, or random sequences, we may simply refer to objects as *randoms*.

A set A is called *left-r.e.*¹ if there exists a uniformly recursive approximation A_0, A_1, A_2, \dots to A such that $A_s \leq_{\text{lex}} A_{s+1}$ for all s . Here $A_s \leq_{\text{lex}} A_{s+1}$ means that either

¹Our definition is analogous to the usual definition of *left-r.e.* for reals which requires that the real admits a recursive approximation from below. In more detail, a real number $X \in [0, 1]$ is called *left-r.e.* if it can be

$A_{s+1} = A_s$ or the least element x of the symmetric difference satisfies $x \in A_{s+1}$. Left-r.e. sets are limit-recursive sets with recursive approximations of a special form. We call A_0, A_1, A_2, \dots a *left-r.e. approximation* of A . Every r.e. set is left-r.e. as $A_s \subseteq A_{s+1}$ implies $A_s \leq_{\text{lex}} A_{s+1}$. Zvonkin and Levin [Zvonkin and Levin 1970] and later Chaitin [Chaitin 1987] showed that there exists a left-r.e. Martin-Löf random set. (Like Chaitin we will fix one and call it Ω .) It follows that each of the classes in Theorem 1.3 contains a left-r.e. member.

A *numbering* φ is a partial-recursive (p.r.) function $\langle e, x \rangle \mapsto \varphi_e(x)$. A numbering φ is a programming language, and φ_e is the e^{th} program in that language. While φ enumerates p.r. functions, our main focus in this paper will be enumerations of sets and reals which admit recursive approximations from below.

Definition 2.1. Let \mathcal{C} be a class of left-r.e. sets². A *left-r.e. numbering* α of \mathcal{C} is a function with range \mathcal{C} given by

$$e \mapsto \lim_{s \rightarrow \infty} \alpha_{e,s} = \alpha_e$$

where:

- (I) $\alpha_{e,s}$ is uniformly recursive in e and s , and
- (II) $\alpha_{e,0}, \alpha_{e,1}, \alpha_{e,2}, \dots$ is a left-r.e. recursive approximation of α_e .

The following definition is a terse review of the arithmetic hierarchy. For a more in-depth discussion see [Soare 1987].

A set $A \subseteq \omega$ is called a Σ_n set if it is Σ_n^0 in the usual sense of recursion theory. The complement of a Σ_n set is a Π_n set. We say that a set A *many-to-one* reduces to a set B , or $A \leq_m B$, if there exists a recursive function f such that for all x , $x \in A \iff f(x) \in B$. A set A is called Σ_n -hard (respectively, Π_n -hard) if for every Σ_n (resp. Π_n) set X , $X \leq_m A$. A set A is Σ_n (resp. Π_n) *complete* if A is a Σ_n (resp. Π_n) set and A is Σ_n -hard (resp. Π_n -hard). The *index set* for a class \mathcal{C} with respect to a (left-r.e.) numbering α is $\{e : \alpha_e \in \mathcal{C}\}$.

We make use of the following classical theorem, and we will prove an analogue for left-r.e. index sets in Theorem 3.7.

THEOREM 2.2 (Σ_3 -REPRESENTATION THEOREM [SOARE 1987]). *Let W_0, W_1, W_2, \dots be an acceptable universal r.e. numbering, and let A be a Σ_3 -set. Then there exists a recursive function f such that for all x ,*

$$\begin{aligned} x \in A &\implies (\forall^\infty y) [W_{f(x,y)} = \omega]; \\ x \notin A &\implies (\forall y) [W_{f(x,y)} \text{ is finite}]. \end{aligned}$$

A left-r.e. numbering of all left-r.e. sets is called *universal*. Similarly, an *r.e. numbering* of a class \mathcal{C} is a mapping $e \mapsto \text{dom } \varphi_e$ for some numbering φ , and an r.e. numbering is *universal* if every r.e. set appears in its range. Universal r.e. numberings are

written in the form

$$X = \sum_{x \in \text{dom } \varphi} 2^{-|x|}.$$

for some numbering φ .

² For reals, the definition of left-r.e. numbering would be similar but, as we see from Remark 2.4, not equivalent. A *left-r.e. numbering* of a class of left-r.e. reals \mathcal{C} is a function with range \mathcal{C} given by

$$e \mapsto \sum_{\sigma \in \text{dom } \varphi_e} 2^{-|\sigma|} \quad (2)$$

for some numbering φ .

known to exist, see [Soare 1987, Definition 4.1]. Universal left-r.e. numberings also exist [Brodhead and Kjos-Hanssen 2009]: if φ_e induces a universal r.e. numbering, then φ_e induces a universal left-r.e. numbering.

We shall use capital letters to denote sequences and sets, but we reserve the capital letter W for r.e. numberings. Greek letters σ and τ will denote finite binary strings, φ and ψ will denote numberings, and α , β , γ , and ζ will be left-r.e. numberings (with an exception in Theorem 2.5).

The following result illustrates a crucial difference between left-r.e. reals and left-r.e. sets:

PROPOSITION 2.3. *The co-infinite left-r.e. sets do not have a left-r.e. numbering.*

PROOF. Suppose that such a numbering α exists, let W_0, W_1, W_2, \dots be a universal r.e. numbering with $W_{d,0}, W_{d,1}, W_{d,2}, \dots$ a recursive approximation of W_d . Then W_d is co-infinite if and only if $W_d = \alpha_e$ for some e , that is:

$$(\exists e) (\forall s, x) (\exists t > s) [\alpha_{e,t}(x) = W_{d,t}(x)].$$

Thus $\{d : W_d \text{ is co-infinite}\}$ is Σ_3 , contradicting the fact that this set is also Π_3 -complete [Soare 1987, Corollary 3.5]. \square

Remark 2.4. On the other hand, every real belongs to the equivalence class of some co-infinite set because every dyadic rational can be represented using infinitely many 0's and finitely many 1's, and every non-dyadic rational can be represented using infinitely many 0's and infinitely many 1's. Brodhead and Kjos-Hanssen [Brodhead and Kjos-Hanssen 2009] showed that there exists a left-r.e. numbering of all left-r.e. sets, hence the class in Proposition 2.3 has a left-r.e. numbering when viewed as real numbers, contrary to the corresponding result for sets.

Theorem 2.5 more precisely describe the relationship between enumerations of left-r.e. sets and left-r.e. reals. A [left-r.e. or r.e.] numbering is called a [left-r.e. or r.e.] *one-one* numbering or left-r.e. *Friedberg* numbering if every member in its range has a unique index.

THEOREM 2.5. *A set C of nonzero reals between 0 and 1 has a left-r.e. numbering α (in the sense of Footnote 2) iff the class of sets*

$$\{A : A \text{ is infinite and } \text{real}(A) \in C\}$$

has a left-r.e. numbering. The same holds for left-r.e. one-one numberings.

PROOF. \implies : Let $\alpha_0, \alpha_1, \dots$ be a (one-one) enumeration with dyadic approximations $\alpha_{e,s}$ to α , let

$$A_{e,s} = \text{set}[(1 - 3^{-s}) \cdot \alpha_{e,s}],$$

and let $A_e = \lim_s A_{e,s}$. Since $\text{real}(A_e) = \alpha_e$ for all e , it remains only to show that A_e is infinite. If α_e is non-dyadic, then A_e is the unique infinite set with $\text{real}(A_e) = \alpha_e$. Otherwise α_e is dyadic, in which case all the sets $A_{e,s}$ are lexicographically less than A_e and so A_e is co-finite. Finally, the numbering A is one-one whenever the numbering α is.

\Leftarrow : If A_0, A_1, A_2, \dots is a list of infinite r.e. sets then the reals

$$\alpha_{e,s} = \sum_{\{x < s : x \in A_{e,s}\}} 2^{-x-1} \cdot A_{e,s}(x)$$

approximate uniformly in e the numbers $\text{real}(A_e)$ from below. Again if the numbering A is one-one then so is α . \square

Garden variety numberings in recursion theory satisfy the s - m - n Theorem [Soare 1987] and are called *acceptable* numberings:

Definition 2.6. A (left-r.e.) numbering φ is called a (left-r.e.) *Gödel numbering* or *acceptable (left-r.e.) numbering* if for every (left-r.e.) numbering ψ there exists a recursive function f such that $\varphi_{f(e)} = \psi_e$ for all e .

Intuitively, the function f in Definition 2.6 translates code from program ψ into program φ . Thus acceptable numberings are maximal: any given numbering can be uniformly translated into any acceptable one. Furthermore, any two acceptable numberings are isomorphic in the sense of [Rogers 1958]. These two properties make the notion of an acceptable numbering rather robust. Moreover, the existence of an acceptable numbering is in a sense equivalent to Church's Thesis via the s - m - n Theorem [Soare 1987].

We show that there is no canonical way to number random sets via acceptable left-r.e. numberings. The class of left-r.e. random reals is a natural example of a class which has a left-r.e. numbering but no maximal (i.e. acceptable) numbering.

Definition 2.7. Let $\mathcal{C} \subseteq \{0, 1\}^\omega$. A set X is a *shift-persistent element* of \mathcal{C} if $\sigma \cap X \in \mathcal{C}$ for every string σ .

THEOREM 2.8. Assume that a family \mathcal{C} has a shift-persistent element and there exists an infinite left-r.e. set $R <_{\text{lex}} \omega$ with $R \notin \mathcal{C}$. Then \mathcal{C} does not have an acceptable left-r.e. numbering.

PROOF. Let X be a shift-persistent member of \mathcal{C} , let R be the missed out infinite set with $R <_{\text{lex}} \omega$, and let $\sigma_0, \sigma_1, \sigma_2, \dots$ be a left-r.e. approximation of R such that all n satisfy $\sigma_n 1111 \dots <_{\text{lex}} \sigma_{n+1} 0000 \dots <_{\text{lex}} R$. Every infinite left-r.e. set has such an approximation. Suppose α is an acceptable left-r.e. numbering of \mathcal{C} .

Fix a left-r.e. approximation $\Omega_0, \Omega_1, \Omega_2, \dots$ for Ω , and let $c_\Omega(n)$ be the first stage for which this approximation has settled on the first n positions. Note that c_Ω dominates every recursive function, otherwise we would infinitely often have $K(\Omega \upharpoonright n) \leq \log n + k$ for some constant k . Now there is a \emptyset' -recursive function F such that $F(n)$ is the first m such that the first m bits of R differ from the first m bits of every α_k with $k \leq c_\Omega(n)$. This function F has an approximation F_s and now one takes the set $\beta_n = \sigma_s \cap X$ for the first stage s such that for all $t \geq s$ it holds that $F_t(n) = F_s(n)$ and the first $F_s(n)$ bits of σ_t exist and are equal to those of σ_s . Note that this σ_s can be found as the function values $F_t(n)$ converge to $F(n)$ and similarly the σ_t converge to R .

Each set β_n is in the list $\alpha_0, \alpha_1, \alpha_2, \dots$ by definition of X . Furthermore, β_n coincides with R on its first $F(n)$ bits while every α_k with $k \leq c_\Omega(n)$ differs from R on its first $F(n)$ bits. Hence $\beta_n \notin \{\alpha_0, \alpha_1, \dots, \alpha_{c_\Omega(n)}\}$. It follows that there is no recursive function f with $\beta_n = \alpha_{f(n)}$ for all n as c_Ω would dominate f . Thus the numbering α cannot be an acceptable numbering of the left-r.e. sets of its type. \square

It follows that there is no canonical way to enumerate random reals:

COROLLARY 2.9. There is no acceptable left-r.e. numbering of either the left-r.e. randoms or the left-r.e. non-randoms (under any reasonable definition of random).

3. ARITHMETIC CLASSIFICATION VIA NUMBERINGS

Unlike r.e. numberings, the existence of left-r.e. numberings admits a neat characterization in terms of Σ_3 sets. As a corollary, we will get that the left-r.e. Martin-Löf random reals are enumerable but not co-enumerable. In order to make concatenation easier, we introduce the following operator on finite strings.

Definition 3.1. For any finite binary string σ , σ_{\downarrow} denotes the string σ with the maximum 1 changed to a 0 (if it exists). If σ consists of all zeros, then $\sigma_{\downarrow} = \sigma$.

A refinement of the following result appears in [Nies 2009, Theorem 3.5.21] using an alternate proof.

LEMMA 3.2 (NIES [2009]). *Let X be a sequence which infinitely often has a prefix of length n followed by $n \cdot 2^n$ zeros. Then X is not Schnorr random.*

PROOF. We exhibit a martingale which Schnorr-succeeds on X . The betting strategy is as follows. For simplicity, let us assume that we start with \$3. For the initial bet, place \$1 on the “1” outcome. Now suppose we have already seen a string σ of length n . If the last digit of σ is “0,” then bet 2^{-n} dollars on the “1” outcome. Otherwise, make the same bet that was made the last time.

We claim this martingale succeeds on X . The martingale loses at most 2^{-n} dollars from betting on the $(n+1)^{\text{st}}$ digit of X . Thus the total money lost from playing over an infinite amount of time is at most \$2. On the other hand, we are bound to eventually reach a string of consecutive zeros of length $n \cdot 2^n$ immediately following $X \upharpoonright n$. At this point, 2^{-n} dollars will be wagered $n \cdot 2^n$ times in a row, for a net gain of \$ n over the interval of zeros. By assumption on X we reach such points infinitely often, and therefore the winnings go to infinity.

Finally we exhibit a recursive function which infinitely often is a correct lower bound for the gambler’s capital. Define a recursive function which guesses at each position that we are at the end of an interval of $n \cdot 2^n$ zeros. The function always outputs n where n is the length of the corresponding interval that would have preceded the long string of zeros. if no such integer n exists, then output 0. Infinitely often this guess will be correct and, as noted in the previous paragraph, we will indeed have at least n dollars at this point. \square

Since weakly 1-generic sets are Kurtz random (Theorem 1.5), Proposition 3.3 below implies that Lemma 3.2 does not carry over for Kurtz random sequences.

PROPOSITION 3.3. *Let X be weakly 1-generic sequence and let f be a recursive function. Then for infinitely many n , $(X \upharpoonright n) \cap 0^{f(n)}$ is a prefix of X .*

PROOF. Let

$$A_n = \{\sigma \cap 0^{f(|\sigma|)} : |\sigma| \geq n\}.$$

For all n , some member of A_n is a prefix of X since A_n is a dense r.e. set. Suppose there are only finitely many prefixes of X of the form $(X \upharpoonright n) \cap 0^{f(n)}$, and let k be greater than the length of the longest such prefix. Then some member of A_k must also be a prefix of X , contradicting the definition of k . \square

Definition 3.4. Let $\sigma_0, \sigma_1, \sigma_2, \dots$ be a sequence of strings where $\sigma_{e,s}$ is a stage s approximation of σ_e . We will say that σ_e *blows up to infinity* if $\lim_s |\sigma_{e,s}| = \infty$, and σ_e *gets kicked to infinity* if σ_j blows up to infinity for some $j < e$.

THEOREM 3.5. *Let $A \subseteq \omega$ be a Σ_3 -set, and let α be an acceptable universal left-r.e. numbering. Then there exist a recursive function g such that*

$$x \in A \implies \alpha_{g(x)} \text{ is Martin-L\"of random};$$

$$x \notin A \implies \alpha_{g(x)} \text{ is not Schnorr random}.$$

PROOF. Let W be an acceptable universal r.e. numbering. Without loss of generality, assume that for all e at most one element of e enters W_e at each stage of its enumeration $\{W_{e,s}\}$ and furthermore at least one W_e increases at each stage. By the

Σ_3 -Representation Theorem, there exists a function f satisfying:

$$\begin{aligned} x \in A &\implies W_{f(x,n)} \text{ is infinite for some } n; \\ x \notin A &\implies W_{f(x,n)} \text{ is finite for all } n. \end{aligned}$$

For each x and s , let

$$\sigma_{0,s}^x = \Omega_s \upharpoonright |W_{f(x,0),s}|,$$

let

$$\begin{aligned} m(e, s) &= \text{greatest stage } t + 1 < s \text{ such that} \\ &\quad \max\{x : \Omega_{e,t+1}(x) = 1\} \neq \max\{x : \Omega_{e,t}(x) = 1\}, \end{aligned}$$

and inductively define

$$\sigma_{n+1,s}^x = 1^{(|\sigma_{n,s}^x|+2) \cdot 2^{|\sigma_{n,s}^x|}} \frown (\Omega_s \upharpoonright |W_{f(x,n+1),m[f(x,n+1),s]}|) \downharpoonright. \quad (3)$$

Roughly speaking, $\sigma_{n+1,s}^x$ consists of a long string of 1's followed by an approximation of Ω . Define the recursive function g by

$$\alpha_{g(x)} = \lim_s \sigma_{0,s}^x \frown \sigma_{1,s}^x \frown \sigma_{2,s}^x \dots \quad (4)$$

By Lemma 3.2, there are enough 1's that if all the σ_n^x 's remain finite, then (4) is not Schnorr random. On the other hand, if some σ_n^x does blow up to infinity, then (4) becomes the Martin-Löf random Ω with some finite prefix attached.

We verify that the approximation in (4) is left-r.e. by analyzing the change between stages s and $s + 1$. By induction, the length of $\sigma_{n,t}^x$ is increasing in t for every n . Let e be the least index such that $\sigma_{e,s+1}^x$ is longer than $\sigma_{e,s}^x$. By minimality, the prefix of 1's at the beginning of this string must remain unchanged but the approximation to Ω increases. In particular,

$$|W_{f(x,e),m[f(x,e),s]}| \neq |W_{f(x,e),m[f(x,e),s+1]}|.$$

Due to the \downharpoonright operator, the 0 at some existing position changes to a 1 in stage $s + 1$. Hence σ_e^x can expand in stage $s + 1$ while permitting a left-r.e. approximation for (4). Finally, the limit in (4) exists because the sequence of reals is increasing and bounded from above.

Suppose that $W_{f(x,n)}$ is infinite for some n , and let e be the least such index. By minimality, $\sigma_j^x = \lim_s \sigma_{j,s}^x$ is finite for all $j < e$. Hence for $e > 0$,

$$\alpha_{g(x)} = \sigma_0^x \frown \sigma_1^x \frown \sigma_2^x \frown \dots \frown 1^{(|\sigma_e^x|+2) \cdot 2^{|\sigma_e^x|}} \frown \Omega,$$

which is Martin-Löf random. All σ_n^x with $n > e$ gets kicked to infinity. The case $e = 0$ is similar.

On the other hand, suppose that $W_{f(x,n)}$ is finite for all n . In this case σ_0^x is finite, and

$$\sigma_{n+1}^x = 1^{(|\sigma_n^x|+2) \cdot 2^{|\sigma_n^x|}} \frown (\Omega_{s_n} \upharpoonright |W_{f(x,n+1),m[f(x,n+1),s_n]}|) \downharpoonright,$$

where s_n is the final stage where $W_{f(x,n+1)}$ increases. Thus infinitely often $\alpha_{g(x)}$ has a prefix of length $|\sigma|$ followed by $(|\sigma| + 2) \cdot 2^{|\sigma|}$ 1's. By Lemma 3.2, $\alpha_{g(x)}$ is not Schnorr random. \square

COROLLARY 3.6. *In any acceptable universal left-r.e. numbering, the indices of the left-r.e. Martin-Löf randoms are Σ_3 -hard.*

Recall that a left-r.e. numbering is called a left-r.e. *Friedberg* numbering if every member in its range has a unique index. Friedberg initiated the study of these numberings in 1958 when he showed that the r.e. sets can be enumerated without repetition [Friedberg 1958]. More recently Kummer [Kummer 1990] gave a simplified proof of Friedberg's result, and Brodhead and Kjos-Hanssen [Brodhead and Kjos-Hanssen 2009] adapted his idea to show that there exists a left-r.e. Friedberg numbering of the left-r.e. Martin-Löf random sets. We now show that left-r.e. Friedberg numberings can be used to characterize Σ_3 -index sets.

THEOREM 3.7. *Let \mathcal{C} be a class of infinite left-r.e. reals which contains a shift-persistent element. Then for any universal left-r.e. numbering α , the following are equivalent:*

- (I) $\{e : \alpha_e \in \mathcal{C}\}$ is a Σ_3 -set.
- (II) There exists a left-r.e. numbering of \mathcal{C} .
- (III) There exists a left-r.e. Friedberg numbering of \mathcal{C} .

PROOF. Let α be any universal left-r.e. numbering, and let

$$\mathcal{C}_\alpha = \{e : \alpha_e \in \mathcal{C}\}.$$

PROOF (I) \iff (II). Suppose that β is a left-r.e. numbering for \mathcal{C} . Then

$$\begin{aligned} \mathcal{C}_\alpha &= \{e : (\exists d) [\alpha_e = \beta_d]\} \\ &= \{e : (\exists d) (\forall n, s) (\exists t > s) [\alpha_{e,t} \upharpoonright n = \beta_{d,t} \upharpoonright n]\}, \end{aligned}$$

so \mathcal{C}_α is a Σ_3 set.

Conversely, assume that $\mathcal{C}_\alpha \in \Sigma_3$ and let γ be an acceptable universal left-r.e. numbering. By Theorem 3.5, there exists a recursive function g such that

$$e \in \mathcal{C}_\alpha \iff \gamma_{g(e)} \text{ is Martin-Löf random.}$$

For sets X , let

$$r_b(X) = \max\{n : (\forall m \leq n) [K(X \upharpoonright m) \geq m - b]\},$$

and in case X has a recursive approximation X_0, X_1, X_2, \dots , then we define a monotonic approximation to r_b as follows:

$$r_{b,s+1}(X) = \max\{r_{b,s}(X), \max\{n : (\forall m \leq n) [K_s(X_s \upharpoonright m) \geq m - b]\}\},$$

where K_s is a monotonically decreasing computable approximation to K . It may not be the case that $\lim_s r_{b,s}(X) = r_b(X)$, however we do achieve $\lim_s r_{b,s}(X) = \infty \iff r_b(X) = \infty$.

Without loss of generality, assume that $\alpha_{e,s}$ has finitely many 1's at each stage s of the recursive approximation. Let C be a shift-persistent element of \mathcal{C} , and let C_0, C_1, C_2, \dots be a left-r.e. approximation for C . Since we want to avoid dealing with α_e 's which are equal to 0, let

$$f(e) = e^{\text{th}} \alpha\text{-index found to be nonzero},$$

and let $t(e)$ be the first stage at which $\alpha_{f(e)}$ appears to be nonzero. For notational convenience, let

$$q(e) = \min\{x : \alpha_{f(e),t(e)}(x) = 1\},$$

and let

$$\xi_{(e,b),s} = \begin{cases} 0^{q(e)} & \text{if } |r_{b,s}(\gamma_{g[f(e)]})| \leq q(e); \\ \alpha_{f(e),s} \upharpoonright r_{b,s}(\gamma_{g[f(e)]}) & \text{otherwise} \end{cases}$$

be the prefix of $\alpha_{f(e),s}$ that has the length of $\gamma_{g[f(e)]}$'s prefix which looks random at stage s . Let

$m(e, s) = \text{greatest stage } t + 1 < s \text{ such that}$

$$\max\{x : \alpha_{f(e),t+1}(x) = 1\} \neq \max\{x : \alpha_{f(e),t}(x) = 1\}.$$

Define a further left-r.e. numbering β by

$$\beta_{\langle e,b \rangle, s+1} = \xi_{\langle e,b \rangle, m(e,s)} \smallfrown C_{s+1}. \quad (5)$$

The operator \smallfrown in (5) is needed to ensure that β is a left-r.e. numbering: whenever $r_{b,m(e,s+1)}(\gamma_{g[f(e)]}) \neq r_{b,m(e,s)}(\gamma_{g[f(e)]})$, this expansion is handled by replacing a “0” with “1” which clears the higher indices, making room for C_{s+1} .

Finally, β_0, β_1, \dots is a left-r.e. numbering for \mathcal{C} . Indeed,

$$\begin{aligned} f(e) \in \mathcal{C}_\alpha &\implies (\exists b) [\gamma_{g[f(e)]} \text{ is Martin-L\"of random with constant } b] \\ &\implies \beta_{\langle e,b \rangle} = \alpha_{f(e)}. \end{aligned}$$

Of course a β -index for the real 0 can be added if necessary. In the case where $\gamma_{g[f(e)]}$ is not Martin-L\"of random with constant b , C_s does not get kicked to infinity but then $\beta_{\langle e,b \rangle} \in \mathcal{C}$ because C is a shift-persistent member of \mathcal{C} . \square

PROOF (II) \iff (III). Assume that \mathcal{C} has a numbering γ . Let C be a shift-persistent element of \mathcal{C} , and let

$$\mathcal{B} = \{1^n \smallfrown C : n \in \omega\} \cup \{X \in \mathcal{C} : X \leq C\}$$

be a subclass of \mathcal{C} . \mathcal{B} is the union of two classes which have left-r.e. numberings and therefore has itself a left-r.e. numbering. A numbering for the latter class is achieved by pausing the enumeration of X whenever it tries to exceed C . Let β be a left-r.e. numbering for \mathcal{B} .

Note that

$$\mathcal{A} := \{X : X \in \mathcal{C} - \mathcal{B}\} = \{X \in \mathcal{C} : (\exists n) [1^n \smallfrown C < X < 1^{n+1} \smallfrown C]\}$$

has a left-r.e. numbering α given by: $\alpha_{\langle e,n,k \rangle, s} =$

$$\begin{cases} 1^n \smallfrown C_s + 2^{-k} & \text{if } (\gamma_{e,s} \upharpoonright k) \smallfrown 0 \leq_{\text{lex}} (1^n \smallfrown C_s \upharpoonright k) \smallfrown 0; \\ \gamma_{e,s} & \text{if } (1^n \smallfrown C_s \upharpoonright k) \smallfrown 0 <_{\text{lex}} (\gamma_{e,s} \upharpoonright k) \smallfrown 0 <_{\text{lex}} (1^{n+1} \smallfrown C_s \upharpoonright k) \smallfrown 0; \\ 1^{n+1} \smallfrown C_s - 2^{-k} & \text{if } (1^{n+1} \smallfrown C_s \upharpoonright k) \smallfrown 0 \leq_{\text{lex}} (\gamma_{e,s} \upharpoonright k) \smallfrown 0. \end{cases}$$

where the triple $\langle e, n, k \rangle$ ranges over values k which are greater than or equal to the index of the least 0 in $1^n \smallfrown C$. The numbering α exploits the fact that if $X \neq 1^n \smallfrown C$, then X and $1^n \smallfrown C$ must differ on some prefix. Strictly speaking, every *tail* of C must be a shift-persistent element in order that each α -index yields a member of \mathcal{C} . Since every member of \mathcal{C} is infinite, however, we can overcome this shortcoming by modifying the tails for $\alpha_{\langle e,n,k \rangle, s}$ to be C_s in the first and third cases.

Using α and β , we now exhibit a Friedberg numbering ζ for $\mathcal{A} \cup \mathcal{B} = \mathcal{C}$. Let

$$M = \{e : (\forall j < e) [\alpha_j \neq \alpha_e]\}.$$

Every member of \mathcal{A} has a unique index in M . Since M is a Σ_2 -set, there exists a \emptyset' -recursive function m whose domain is M . Let m_0, m_1, m_2, \dots be a recursive approximation to m . Using this approximation, we shall design ζ in such a way that each α -indexed real in M occurs at exactly one ζ -index, and the remaining ζ -indices will be home to the β -indexed reals.

We define a function $f : \omega \mapsto (\omega \cup \{\infty\}) \times \{\alpha, *\}$ which maps ζ -indices to either α -indices or $*$'s. The ∞ symbol is used for destroyed β indices which are (or never were)

attached to ζ -indices, and the α and $*$ symbols indicate whether the particular ζ -index is following an α -index or a β -index. If $f(e) = \langle x, * \rangle$ for some x , $f(e)$ “explodes” and we say that the ζ -index e has been *destroyed*. $f_s : \omega \mapsto ((\omega \cup \{\infty\}) \times \{\alpha, *\}) \cup \{\uparrow\}$ will be a recursive approximation to f based on the recursive approximation m_s . ζ -indices that are destroyed at some stage take on β -indices in the limit (rather than α -indices). We shall also keep track of which β -indices have been taken on by ζ -indices: G_s will be the set of β -indices which have been ζ -used by stage s . We will achieve $\lim G_s = \omega$. Since \mathcal{C} contains only infinite sets, every α -indexed real is less than some β -indexed real, and therefore we can use β as a garbage can to collect for those approximations $m_s(e)$ which turned out to be wrong. We shall also have an auxiliary recursive function $r(s)$ which marks the boundary between the ζ -indices which are following values in $\omega \cup \{*\}$ and those whose value is \uparrow at stage s .

The construction is as follows:

. *Stage 0.*

Set $G_0 = \emptyset$, $r(0) = 0$, $f_0(e) = \uparrow$, and $\zeta_{e,0} = 0$ for all $e \geq 0$.

. *Stage $s + 1$.* Let

$$\begin{aligned} A &= \{x < s : m_{s+1}(x) \uparrow \text{ and } m_s(x) \downarrow\}, \\ X &= \{x < s : m_{s+1}(x) \downarrow \text{ and } m_s(x) \uparrow\}, \end{aligned}$$

let $\{a_1, a_2, \dots, a_k\}$ be the indices below or equal to $r(s)$ satisfying $f_s(e_i) \in A$, and let $\{x_1, x_2, \dots, x_d\}$ be the indices below or equal to $r(s)$ satisfying $f_s(e_i) \in X$. We destroy all followers of $\{x_1, \dots, x_d\}$, and create new followers for $\{a_1, \dots, a_k\}$:

$$f_{s+1}(n) = \begin{cases} \langle x_i, * \rangle & \text{if } f_s(n) = \langle x_i, \alpha \rangle \text{ for some } 1 \leq i \leq d; \\ \langle a_i, \alpha \rangle & \text{if } n = r(s) + i \text{ for some } 1 \leq i \leq k; \\ \langle s, \alpha \rangle & \text{if } n = r(s) + k + 1; \\ \langle \infty, * \rangle & \text{if } n = r(s) + k + 2; \\ f_s(n) & \text{otherwise.} \end{cases} \quad (6)$$

The ζ -index $r(s) + k + 1$ is used to introduce a new α -index, and the ζ -index $r(s) + k + 2$ is used to ensure that some new β -index is taken up at this stage. Set $r(s + 1) = r(s) + k + 2$.

Next, assign new reals from \mathcal{B} to the ζ -indices that were destroyed in this stage.

— Let

$$y_1 = (\mu n) [\beta_{n,s} > \zeta_{x_1,s} \quad \& \quad n \notin G_s]$$

and inductively for $0 \leq i \leq d$,

$$y_{i+1} = (\mu n) [\beta_{n,s} > \max\{\zeta_{x_{i+1},s}, \beta_{y_i}\} \quad \& \quad n \notin G_s].$$

Choose the least β -index not yet assigned to a ζ -index and call it z :

$$z = \min\{n : n \notin \{y_1, y_2, \dots, y_d\} \text{ and } n \notin G_s\}. \quad (7)$$

This choice of z ensures that every member of \mathcal{B} will have some index in ζ .

— Set

$$\zeta_{n,t} = \begin{cases} \beta_{y_i,t} & \text{if } f_{s+1}(n) = \langle x_i, * \rangle \text{ for some } 1 \leq i \leq d; \\ \beta_{z,t} & \text{if } n = r(s + 1). \end{cases}$$

for all $t > s$.

— Set $G_{s+1} = G_s \cup \{y_0, y_1, \dots, y_k, z\}$.

For the remaining ζ -indices which have not been destroyed in this stage or some previous stage, continue following α -indices:

$$\zeta_{e,s+1} = \begin{cases} 0 & \text{if } f_{s+1}(e) = \uparrow; \\ \alpha_{f_{s+1}(e),s+1} & \text{if } f_{s+1}(e) \notin \{\langle n, * \rangle : n \in \omega\} \cup \{\uparrow\}. \end{cases} \quad (8)$$

By induction on stages, (6) and (8) ensure that for all s and $e \leq s$, there exists a unique n such that

$$\zeta_{n,s+1} = \alpha_{\text{first projection of } f_{s+1}(e), s+1}.$$

Since each sequence $\{\alpha_{\text{first projection of } f_{s+1}(e)}\}$ converges to a unique member in the range of α on the set of indices $e \in \text{dom } m$, it follows that there is a unique ζ -index for each real in \mathcal{A} . Indeed for $e \notin \text{dom } m$, the approximation for $m(e)$ may oscillate between convergence and divergence infinitely often, but we simply introduce a fresh ζ -index for an unused member of \mathcal{B} each time this happens and therefore α_e will not occupy a ζ -index in the limit. Furthermore (6) and (7) ensure that there is a unique ζ -index for each real in \mathcal{B} .

Finally, $\zeta_e \in \mathcal{A} \cup \mathcal{B}$ for all e . If the index e is destroyed at some stage in the construction, then some β -index n is assigned at that stage and $\zeta_e = \beta_n$. On the other hand if index e is never destroyed, then ζ_e takes an α -index, namely $\zeta_e = \alpha_{\text{first projection of } f(e)}$. \square

Hence (I) \iff (II) \iff (III).

COROLLARY 3.8. *The following classes have left-r.e. numberings:*

- (I) *the left-r.e. Martin-Löf random sets,*
- (II) *the left-r.e. Kurtz non-random sets, and*
- (III) *the infinite left-r.e. sets.*
- (IV) *the infinite r.e. sets.*

Proposition 3.9 below contrasts with Corollary 3.8(IV). This dichotomy does not surprise us too much as the recursive sets are also enumerable if viewed as r.e. characteristic functions, what is well-known to be impossible for recursive functions. To see such an enumeration, we start with an enumeration of the binary p.r. functions, f_0, f_1, f_2, \dots . We can uniformly interpret each f_i as the recursive set whose characteristic function is the truncation of f_i up to the highest number n such that $f_i(x) \downarrow$ for all $x < n$, followed by the constant zero function. Then the indices for total functions will yield the characteristic functions for the recursive sets, and the non-total functions will yield finite sets which are also recursive.

PROPOSITION 3.9. *There is no r.e. numbering of the infinite r.e. sets.*

PROOF. Suppose that A_0, A_1, A_2, \dots were an r.e. numbering of the infinite r.e. sets. Search for an $a_0 \in A_0$, and let $b_0 = a_0 + 1$. Next, search for an $a_1 \in A_1$ which is greater than b_0 , and let $b_1 = a_1 + 1$. Continuing the diagonalization, find $a_2 \in A_2$ which is greater than b_1 and let $b_2 = a_2 + 1$, and proceed similarly for b_3, b_4, \dots . Now $\{b_0 < b_1 < b_2 < \dots\}$ is an infinite r.e. set which disagrees from the n^{th} r.e. set at a_n . \square

It remains to show that the hypothesis “contains a shift-persistent element” is necessary in Theorem 3.7.

THEOREM 3.10. *There exists a Σ_3 -class of infinite left-r.e. reals which contains no shift-persistent element and has no left-r.e. numbering.*

PROOF. Let α be a universal left-r.e. numbering and define the following α -index set:

$$X = \{e : (\exists x) [x \notin \alpha_e \cup \Omega] \text{ and } (\forall y < x) [y \in \alpha_e \iff y \notin \Omega] \text{ and } (\forall y > x) [y \in \alpha_e \iff y \text{ is odd}]\}. \quad (9)$$

By the third line, X is infinite, and by the second line, X contains no shift-persistent element. Furthermore, (9) is a Σ_2 -formula with a \emptyset' -recursive predicate, hence $X \in \Sigma_3$. If X would have a left-r.e. numbering, then by the first line, $\bar{\Omega}$ would be the lexicographic supremum of all the approximations occurring to members of this left-r.e. numbering and $\bar{\Omega}$ would be a left-r.e. set, contradicting that Ω is nonrecursive. \square

Also along the lines of randomness, we note that the class of left-r.e. reals X satisfying $X + \Omega \leq 1$ has a Π_1 index set (in any numbering), has no shift-persistent element, and has no left-r.e. numbering. Indeed if this class had a left-r.e. numbering, then Ω would be recursive.

COROLLARY 3.11. *The left-r.e. Martin-Löf non-random reals, computable non-random reals, and Schnorr non-random reals have no left-r.e. numberings. Hence none of these classes has a Σ_3 index set in any universal left-r.e. numbering.*

PROOF. These classes are Π_3 -hard in any acceptable numbering by Corollary 3.6. It follows from Theorem 3.7 that none of these classes are effectively enumerable and hence cannot be Σ_3 in any universal left-r.e. numbering. \square

4. WEAKLY 1-GENERIC SETS

We examine left-r.e. numberings for Kurtz random, bi-immune, bi-hyperimmune, and weakly 1-generic sets. We introduced weakly 1-generic sets in Definition 1.4.

Definition 4.1. An infinite set is *immune* if it contains no infinite recursive subset. Even stronger, a set $A = \{a_0 < a_1 < \dots\}$ is *hyperimmune* if there exists no recursive function f such that $f(n) > a_n$ for all n . It is *bi-(hyper)immune* if both A and the complement \bar{A} are (hyper)immune.

THEOREM 4.2. *Let $A \subseteq \omega$ be a Σ_3 -set, and let α be an acceptable universal left-r.e. numbering. Then there exist a recursive function g such that*

$$\begin{aligned} x \in A &\implies \alpha_{g(x)} \text{ is co-finite;} \\ x \notin A &\implies \alpha_{g(x)} \text{ is weakly 1-generic.} \end{aligned}$$

PROOF. Let W be an acceptable universal r.e. numbering. By the Σ_3 -Representation Theorem, there exists a recursive function f such that:

$$\begin{aligned} x \in A &\implies W_{f(x,n)} \text{ is infinite for some } n; \\ x \notin A &\implies W_{f(x,n)} \text{ is finite for all } n. \end{aligned}$$

The idea now is make $\alpha_{g(x)}$ a sequence of the form

$$\alpha_{g(x)} = \sigma_0 \frown 0 \frown \sigma_1 \frown 0 \frown \sigma_2 \frown 0 \dots$$

such that $\sigma_0 \frown 0 \frown \sigma_1 \frown 0 \frown \dots \frown \sigma_n$ is a member of W_n whenever W_n is dense and $W_{f(x,e)}$ is finite for all e . If on the other hand $W_{f(x,e)}$ is infinite for some e , then some σ_n will blow up to infinity and $\alpha_{g(x)}$ will be co-finite.

For every n , let

$$\tau_{n,s} = \sigma_{0,s} \frown 0 \frown \sigma_{1,s} \frown 0 \frown \dots \frown \sigma_{n,s}.$$

At Stage 0, $\sigma_{e,0} = 0$ for all e , and $\alpha_{g(x),0} = \sigma_{0,0} \frown 0 \frown \sigma_{1,0} \frown 0 \frown \sigma_{2,0} \frown 0 \frown \dots$. At Stage $s + 1$, let e be the least index, if one exists, such that either:

- (1) $|W_{f(e,x),s+1}| > |W_{f(e,x),s}|$, or
- (2) some member of $W_{e,s+1}$ extends $\tau_e \frown 1$, and no member of $W_{e,s+1}$ is a prefix of τ_e .

If condition 1 is satisfied, let $\sigma_{e,s+1} = \sigma_{e,s} \frown 1$ so that σ_e becomes longer. Otherwise, let $\sigma_{e,s+1}$ be an extension of $\sigma_{e,s} \frown 1$ such that $\tau_{e,s+1} \in W_{e,s+1}$. This latter case aims to make $\alpha_{g(x)}$ weakly 1-generic. In either case, $\sigma_{j,s+1} = \sigma_{j,s}$ for $j \neq e$. If no such e exists, skip to Stage $s + 2$.

The sequence $\{\alpha_{g(x),s}\}$ is indeed a left-r.e. approximation. In each stage s where some action takes place in the construction, the 0 following $\sigma_{e,s}$ is changed to a 1 before this string is extended.

We claim that if $W_{f(x,n)}$ is finite for all n , then $\alpha_{g(x)}$ will be weakly 1-generic. By some stage s , $W_{f(0,x),s}$ must stop expanding. Whether or not W_0 is dense, σ_0 will change at most one time after stage s , and therefore τ_0 settles by some stage t_0 . If W_0 is dense, then τ_0 will contain a member of W_0 . Similarly $W_{f(1,x),s}$ must stop expanding at some point after stage t_0 , τ_1 will eventually contain a prefix of W_1 if W_1 is dense, and $\tau_{1,t}$ settles by some stage t_1 . Continuing by induction, we see that $\alpha_{g(x)}$ is weakly 1-generic.

If $W_{f(x,n)}$ is infinite for some least n , then the argument in the previous paragraph shows that $\tau_{n-1,t}$ eventually settles, and then infinitely often $\sigma_{n,s+1} = \sigma_{n,s} \frown 1$ and so $\alpha_{g(x)}$ is co-finite. \square

Since every weakly 1-generic set is both hyperimmune [Nies 2009, Proposition 1.8.48] and Kurtz random (Theorem 1.5) we have the following:

COROLLARY 4.3. *In any acceptable universal left-r.e. numbering, the index sets for the following classes are Π_3 -hard:*

- (I) *the left-r.e. immune sets,*
- (II) *the left-r.e. hyperimmune sets,*
- (III) *the left-r.e. bi-immune sets,*
- (IV) *the left-r.e. bi-hyperimmune sets,*
- (V) *the left-r.e. weakly 1-generic sets,*
- (VI) *the left-r.e. Kurtz random sets.*

From Theorem 3.7 we also have the following result.

COROLLARY 4.4. *In any universal left-r.e. numbering, the classes listed in Corollary 4.3 have $\Pi_3 - \Sigma_3$ index sets. Moreover there exists a left-r.e. numbering for each of the corresponding complementary classes.*

It is known that every Kurtz random is bi-immune [Kurtz 1981], but the reverse inclusion does not hold [Bienvenu et al. 2010]. We can also separate the left-r.e. versions of these notions.

PROPOSITION 4.5. *There exists a left-r.e. bi-hyperimmune set which is not Kurtz random.*

PROOF. Let A be any bi-hyperimmune left-r.e. set. Then $A \oplus A$ is bi-hyperimmune but not Kurtz random since a recursive martingale can win on every second bit. \square

The reverse of Proposition 4.5 holds as well: Chaitin's Ω is an example of a left-r.e. Martin-Löf random (in particular, Kurtz random) which, by Lemma 3.2, is not hyperimmune (in particular, not bi-hyperimmune).

5. CLASSES OF HIGHER COMPLEXITY

We now investigate the complex randomness notions of Schnorr randomness and computable randomness. As we shall see, neither of these left-r.e. classes, nor their complements, have left-r.e. numberings. A set A is called *high* if $A' \geq_T \emptyset''$. A theorem of Nies, Stephan, and Terwijn [Nies et al. 2005] shows the existence of left-r.e. Schnorr randoms which are not Martin-Löf random:

THEOREM 5.1 (NIES, STEPHAN, TERWIJN [NIES ET AL. 2005]). *The following statements are equivalent for any set A :*

- (I) A is high.
 - (II) There is a set $B \equiv_T A$ which is computably random but not Martin-Löf random.
 - (III) There is a set $C \equiv_T A$ which is Schnorr random but not computably random.
- In the case that A is left-r.e. and high, the sets B and C can be chosen as left-r.e. sets as well.*

Furthermore, Downey and Griffith [Downey and Griffiths 2004; Downey and Hirschfeldt 2010] proved that every left-r.e. Schnorr random real is high. Therefore

FACT 5.2. *A left-r.e. set X is high $\iff X$ Turing equivalent to a left-r.e. Schnorr random set $\iff X$ is Turing equivalent to a left-r.e. computably random set.*

In his PhD thesis [Schwarz 1982], Schwarz characterized the complexity of the high r.e. degrees:

THEOREM 5.3 (SCHWARZ [1982], [SOARE 1987]). *In any acceptable universal r.e. numbering W_0, W_1, W_2, \dots , $\{e : W_e \text{ is high}\}$ is Σ_5 -complete.*

Using this Schwarz's theorem, we obtain the following enumeration result.

THEOREM 5.4. *Let \mathcal{C} be a class of left-r.e. reals such that:*

- (I) *Every member of \mathcal{C} is high, and*
- (II) *every high r.e. set is Turing equivalent to some member of \mathcal{C} .*

Then for any universal left-r.e. numbering α , $\{e : \alpha_e \in \mathcal{C}\}$ is not a Σ_4 -set and hence is neither enumerable nor co-enumerable.

PROOF. Let \mathcal{C} be a class satisfying the hypothesis of the theorem, let W be an acceptable universal r.e. numbering, let Φ denote a Turing functional, and suppose that

$$\alpha_i \in \mathcal{C} \iff (\exists n_1) (\forall n_2) (\exists n_3) (\forall n_4) [P(i, n_1, n_2, n_3, n_4)].$$

for some recursive predicate P .

For convenience assume that whenever a computation $\Phi_j^{W_{e,t}}$ is injured, it is undefined for at least one stage; then

$$\begin{aligned} W_e \text{ is high} &\iff (\exists i, j) \left[\alpha_i \in \mathcal{C} \ \& \ \alpha_i = \Phi_j^{W_e} \right] \\ &\iff (\exists i, j, n_1) (\forall x, n_2) (\exists t) (\exists n_3) (\forall u > t) (\forall n_4) \\ &\quad \left[P(i, n_1, n_2, n_3, n_4) \ \& \ \alpha_{i,u}(x) = \Phi_{j,u}^{W_{e,u}}(x) \right]. \end{aligned}$$

Thus $\{e : W_e \text{ is high}\}$ is a Σ_4 -set, contrary to Theorem 5.3. \square

COROLLARY 5.5. *Neither the Schnorr random sets nor the computably random sets reals are Σ_4 in any universal left-r.e. numbering. Hence neither class nor its complement has a left-r.e. numbering.*

PROOF. By Fact 5.2, the left-r.e. Schnorr random sets and the left-r.e. computably random sets satisfy the hypothesis of the Theorem 5.4. Apply Theorem 3.7. \square

It remains to characterize the hardness of computable random sets and Schnorr random sets in an acceptable universal left-r.e. numbering. For the remainder of this paper, we fix an acceptable universal left-r.e. numbering α and an acceptable universal r.e. numbering W . The *principal function* of a set $A = \{a_0 < a_1 < a_2 < \dots\}$ is given by $n \mapsto a_n$; we write $p_A(n) = a_n$. We will be particularly interested in the principal functions of co-r.e. sets, so we use the abbreviation $p_{\bar{e}}$ for $p_{\overline{W_e}}$. We say that a function $f : \omega \rightarrow \omega$ is *dominating* if it dominates all recursive functions.

THEOREM 5.6. *There is a Turing reduction procedure Φ and a recursive function g such that for all e ,*

- (I) $\Phi^{p_{\bar{e}}}$ is a left-r.e. real,
- (II) $\alpha_{g(e)} = \Phi^{p_{\bar{e}}}$, and
- (III) Φ^f is computably random if f is dominating.

PROOF. This fact follows from the proof of Nies, Stephan, Terwijn [Nies et al. 2005, Theorem 4.2, (I) implies (II), r.e. case] which appears as Theorem 5.1 in this paper. \square

A set A is *low* if $A' \leq_T \emptyset'$, and a function is *low* if it is computable from a low set. A function f is *diagonally non-recursive (DNR)* if for some numbering φ and every e , the value $\varphi_e(e)$, if defined, differs from $f(e)$.

LEMMA 5.7. *A low left-r.e. set cannot compute a Schnorr random.*

PROOF. Suppose such a real A computes a Schnorr random set X . Since X is not high, X must also be Martin-Löf random (by Theorem 5.1). Kučera showed that every Martin-Löf random set computes a DNR function [Kučera 1985], [Kjos-Hanssen et al. 2006, Theorem 6], so A computes a DNR function. Moreover A has r.e. Turing degree because it is truth-table equivalent to the r.e. set $\{\sigma : \sigma \cap 0^\omega \leq A\}$. An r.e. set computes a DNR-function if and only if the set is Turing complete [Arslanov 1981][Jockusch et al. 1989][Kjos-Hanssen et al. 2006, Corollary 9][Kjos-Hanssen et al. 2011], hence $A \equiv_T \emptyset'$. This contradicts the fact that A is low. \square

An r.e. set A is *maximal* if for each r.e. set W with $A \subseteq W$, either $\omega \setminus W$ or $W \setminus A$ is finite. Friedberg [Friedberg 1958] proved that maximal sets exist.

THEOREM 5.8. *For every $A \in \Pi_4$, there exists a recursive function f such that for all e ,*

$$e \in A \implies \alpha_{f(e)} \text{ is computably random;} \quad (10)$$

$$e \notin A \implies \alpha_{f(e)} \text{ is not Schnorr random.} \quad (11)$$

PROOF. Let us fix a Π_4 -complete set A ; By [Soare 1987, XII. Exercise 4.26], there is a recursive function h such that

$$e \in A \iff W_{h(e)} \text{ is maximal} \iff W_{h(e)} \text{ is not low.}$$

Martin and Tennenbaum showed that the principal function of the complement of a maximal set dominates all recursive functions [Soare 1987, XI. Proposition 1.2]. Using this result and the function g given by Theorem 5.6,

$$\begin{aligned} W_{h(e)} \text{ is maximal} &\implies p_{\overline{h(e)}} \text{ is dominating} \\ &\implies \alpha_{g[h(e)]} = \Phi^{p_{\overline{h(e)}}} \text{ is computably random,} \end{aligned}$$

and by Lemma 5.7 with Theorem 5.6(I),

$$\begin{aligned} W_{h(e)} \text{ is not maximal} &\implies p_{\overline{h(e)}} \text{ is low} \\ \implies \alpha_{g[h(e)]} = \Phi^{\overline{p_{h(e)}}} &\text{ is not Schnorr random.} \end{aligned}$$

The function $f = g \circ h$ witnesses the conclusion of this theorem. \square

Note that if we replaced “computably random” with “Martin-Löf random” in (10), we would obtain a characterization of Σ_3 sets rather than Π_4 sets (care of Theorem 3.5). Since every computable random is Schnorr random (Theorem 1.3), we obtained an optimal hardness result:

COROLLARY 5.9. *In any acceptable universal left-r.e. numbering, both the indices of the Schnorr random sets and the indices of the computably random sets are Π_4 -complete.*

We summarize our main results in Table I. A theorem in a forthcoming paper [Stephan and Teutsch Stephan and Teutsch] states that every \emptyset' -recursive 1-generic set has a co-r.e. indifferent set which is retraceable by a recursive function. It follows that for each the families of randoms listed in Table I, there exists a universal left-r.e. numbering which makes the set of the indices for that class 1-generic. Therefore we cannot obtain any arithmetic hardness results for index sets in the general case of universal left-r.e. numberings.

Table I. Calculated complexities

Left-r.e. family	Complexity	Hardness*
Martin-Löf randoms	$\Sigma_3 - \Pi_3$ [3.11]	Σ_3 -hard [3.6]
computable randoms	$\Pi_4 - \Sigma_4$ [5.5]	Π_4 -hard [5.9]
Schnorr randoms	$\Pi_4 - \Sigma_4$ [5.5]	Π_4 -hard [5.9]
Kurtz randoms	$\Pi_3 - \Sigma_3$ [4.4]	Π_3 -hard [4.3]
bi-immune sets	$\Pi_3 - \Sigma_3$ [4.4]	Π_3 -hard [4.3]

Note: Complexities listed hold for any universal left-r.e. numbering.

*Hardness results are for *acceptable* universal left-r.e. numberings.

We can separate most of the adjacent left-r.e. classes in Table I simply by observing differences in arithmetic complexity (and using the well-known result Theorem 1.3). The remaining separations follow from Theorem 5.1 and Proposition 4.5. All of these separations were previously known, with the possible exception of a left-r.e. Kurtz random which is not bi-immune.

Among the families Table I, only the Martin-Löf randoms have a left-r.e. numbering, and among the complementary families only the Kurtz non-randoms and non-bi-immune sets do (by Theorem 3.7).

6. EXPANDING THE VOCABULARY

In Section 2, we identified left-r.e. sets as limit-recursive sets with recursive approximations of a special form. However there are other easy to describe limit-recursive sets which are Martin-Löf random but not left-r.e. For example, $\{x : 2x \in \Omega\}$ is Martin-Löf random and low by van Lambalgen’s Theorem [Downey and Hirschfeldt 2010; van Lambalgen 1987] but not left-r.e. (as left-r.e. Martin-Löf random sets are weak truth-table complete [Nies 2009, Corollary 3.2.31]). See [Jain et al. 2011, Proposition 13] for an elementary explanation why $\{x : 2x \in \Omega\}$ and $\{x : 2x + 1 \in \Omega\}$ cannot both have left-r.e. approximations.

QUESTION 6.1. If $A = a_0 < a_1 < a_2 < \dots$ is an infinite r.e. (co-r.e.) set, and Ω is a left-r.e. Martin-Löf random, is the set

$$\Omega(a_0)\Omega(a_1)\Omega(a_2)\dots \quad (12)$$

Martin-Löf random? If not, which classes of sequences of the form (12) have numberings?

REFERENCES

- ARSLANOV, M. M. 1981. Some generalizations of a fixed-point theorem. *Izv. Vyssh. Uchebn. Zaved. Mat.* 5, 9–16.
- BIENVENU, L., STEPHAN, F., AND TEUTSCH, J. 2010. How powerful are integer-valued martingales? In *Programs, Proofs, Processes (CiE 2010)*. Lecture Notes in Computer Science. Springer-Verlag, Berlin, Heidelberg, 59–68.
- BRODHEAD, P. AND KJOS-HANSSSEN, B. 2009. Numberings and randomness. In *Mathematical Theory and Computational Practice*. Lecture Notes in Computer Science Series, vol. 5635. Springer-Verlag, Berlin, Heidelberg, 49–58.
- CHAITIN, G. J. 1987. Incompleteness theorems for random reals. *Adv. in Appl. Math.* 8, 2, 119–146.
- DOWNEY, R. G. AND GRIFFITHS, E. J. 2004. Schnorr randomness. *J. Symbolic Logic* 69, 2, 533–554.
- DOWNEY, R. G., GRIFFITHS, E. J., AND REID, S. 2004. On Kurtz randomness. *Theoret. Comput. Sci.* 321, 2-3, 249–270.
- DOWNEY, R. G. AND HIRSCHFELDT, D. R. 2010. *Algorithmic randomness and complexity*. Theory and Applications of Computability. Springer, New York.
- FRIEDBERG, R. M. 1958. Three theorems on recursive enumeration. I. Decomposition. II. Maximal set. III. Enumeration without duplication. *J. Symb. Logic* 23, 309–316.
- HITCHCOCK, J. M., LUTZ, J. H., AND TERWIJN, S. A. 2007. The arithmetical complexity of dimension and randomness. *ACM Trans. Comput. Logic* 8, 2, 13.
- JAIN, S., STEPHAN, F., AND TEUTSCH, J. 2011. Closed left-r.e. sets. In *Theory and Applications of Models of Computation (TAMC 2011)*, M. Ogihara and J. Tarui, Eds. Lecture Notes in Computer Science Series, vol. 6648. Springer Berlin / Heidelberg, 218–229.
- JOCKUSCH, JR., C. G., LERMAN, M., SOARE, R. I., AND SOLOVAY, R. M. 1989. Recursively enumerable sets modulo iterated jumps and extensions of Arslanov’s completeness criterion. *J. Symbolic Logic* 54, 4, 1288–1323.
- KJOS-HANSSSEN, B., MERKLE, W., AND STEPHAN, F. 2006. Kolmogorov complexity and the Recursion Theorem. In *STACS 2006*. Lecture Notes in Computer Science. Springer-Verlag, Berlin, Heidelberg, 149–161.
- KJOS-HANSSSEN, B., MERKLE, W., AND STEPHAN, F. 2011. Kolmogorov complexity and the Recursion Theorem. *Transactions of the American Mathematical Society*. Posted on April 27, 2011, PII: S 0002-9947(2011)05306-7 (to appear in print).
- KUČERA, A. 1985. Measure, Π_1^0 -classes and complete extensions of PA. In *Recursion theory week (Oberwolfach, 1984)*. Lecture Notes in Math. Series, vol. 1141. Springer, Berlin, 245–259.
- KUMMER, M. 1990. An easy priority-free proof of a theorem of Friedberg. *Theoret. Comput. Sci.* 74, 2, 249–251.
- KURTZ, S. 1981. Randomness and genericity in the degrees of unsolvability. Ph.D. thesis, University of Illinois at Urbana.
- LEVIN, L. A. 1974. Laws of information conservation (nongrowth) and aspects of the foundation of probability theory. *Problems of Information Transmission* 10, 3, 206–210.

- LI, M. AND VITÁNYI, P. 2008. *An introduction to Kolmogorov complexity and its applications* Third Ed. Texts in Computer Science. Springer, New York.
- MARTIN-LÖF, P. 1966. The definition of random sequences. *Information and Control* 9, 602–619.
- NIES, A. 2009. *Computability and Randomness*. Oxford University Press, Inc., New York, NY, USA.
- NIES, A., STEPHAN, F., AND TERWIJN, S. A. 2005. Randomness, relativization and Turing degrees. *J. Symbolic Logic* 70, 2, 515–535.
- ODIFREDDI, P. G. 1999. *Classical recursion theory. Vol. II*. Studies in Logic and the Foundations of Mathematics. North-Holland Publishing Co., Amsterdam.
- ROGERS, JR., H. 1958. Gödel numberings of partial recursive functions. *J. Symb. Logic* 23, 331–341.
- SCHNORR, C.-P. 1971a. A unified approach to the definition of random sequences. *Math. Systems Theory* 5, 246–258.
- SCHNORR, C.-P. 1971b. *Zufälligkeit und Wahrscheinlichkeit. Eine algorithmische Begründung der Wahrscheinlichkeitstheorie*. Lecture Notes in Mathematics, Vol. 218. Springer-Verlag, Berlin.
- SCHWARZ, S. 1982. Quotient lattices, index sets, and recursive linear orderings. Ph.D. thesis, University of Chicago.
- SOARE, R. I. 1987. *Recursively enumerable sets and degrees*. Perspectives in Mathematical Logic. Springer-Verlag, Berlin. A study of computable functions and computably generated sets.
- STEPHAN, F. AND TEUTSCH, J. Things that can be made into themselves. Manuscript.
- VAN LAMBALGEN, M. 1987. Random sequences. Ph.D. thesis, University of Amsterdam.
- WANG, Y. 1996. Randomness and complexity. Ph.D. thesis, Mathematisch-Naturwissenschaftlichen Gesamtfakultät, Universität Heidelberg.
- ZVONKIN, A. K. AND LEVIN, L. A. 1970. The complexity of finite objects and the development of the concepts of information and randomness by means of the theory of algorithms. *Russian Mathematical Surveys* 25, 6, 83–124.

Received August 2010; revised June 2011; accepted July 2011